arXiv:1001.1881v4 [math.QA] 15 Nov 2010

PERIODICITIES OF T AND Y-SYSTEMS, DILOGARITHM IDENTITIES, AND CLUSTER ALGEBRAS II: TYPES C_r, F_4, AND G_2

REI INOUE, OSAMU IYAMA, BERNHARD KELLER,
ATSUO KUNIBA, AND TOMOKI NAKANISHI

Abstract. We prove the periodicities of the restricted T and Y-systems associated with the quantum affine algebra of type C_r, F_4, and G_2 at any level. We also prove the dilogarithm identities for these Y-systems at any level. Our proof is based on the tropical Y-systems and the categorification of the cluster algebra associated with any skew-symmetric matrix by Plamondon.

Contents

1. Introduction 1
2. Main results 2
3. Type C_r 8
4. Type F_4 21
5. Type G_2 27
6. Mutation equivalence of quivers 36
References 36

1. Introduction

This is the continuation of the paper [IIKKN]. In [IIKKN], we proved the periodicities of the restricted T and Y-systems associated with the quantum affine algebra of type B_r at any level. We also proved the dilogarithm identities for these Y-systems at any level. Our proof was based on the tropical Y-systems and the categorification of the cluster algebra associated with any skew-symmetric matrix by Plamondon [P1, P2]. In this paper, using the same method, we prove the corresponding statements for type C_r, F_4, and G_2, thereby completing all the nonsimply laced types.

The results are basically parallel to type B_r. Since the common method and the proofs of the statements for type B_r were described in [IIKKN] in detail, in this paper, we skip the proofs of most statements, and concentrate on presenting the results with emphasis on the special features of each case. Notably, the tropical Y-system at level 2, which is the core part in the entire method, is quite specific to each case.

While we try to make the paper as self-contained as possible, we also try to minimize duplication with [IIKKN]. Therefore, we have to ask the reader's patience to refer to the companion paper [IIKKN] for the things which are omitted. In

---

2010 *Mathematics Subject Classification.* Primary 13F60; Secondary 17B37.

particular, basic definitions for cluster algebras are summarized in [IIKKN, Section 2.1].

The organization of the paper is as follows. In section 2 we present the main results as well as the T and Y-systems for each type. In Section 3 the results for type C_r are presented. The tropical Y-system at level 2 is the key and described in detail in Section 3.6. In Section 4 the results for type F_4 are presented. In Section 5 the results for type G_2 are presented. In Section 6 we list the known mutation equivalences of quivers corresponding to the T and Y-systems.

2. MAIN RESULTS

2.1. **Restricted T and Y-systems of types C_r, F_4, and G_2.**

Let X_r be the Dynkin diagram of type C_r, F_4, or G_2 with rank r, and $I = \{1, \dots, r\}$ be the enumeration of the vertices of X_r as below.







Let h and $h^\vee$ be the Coxeter number and the dual Coxeter number of X_r, respectively.

(2.1)

X_r	C_r	F_4	G_2
h	$2r$	12	6
$h^\vee$	$r+1$	9	4

We set numbers t and t_a ($a \in I$) by

$$t = \begin{cases} 2 & X_r = C_r,\, F_4 \\ 3 & X_r = G_2, \end{cases} \qquad t_a = \begin{cases} 1 & \alpha_a\text{: long root} \\ t & \alpha_a\text{: short root.} \end{cases} \tag{2.2}$$

For a given integer $\ell \geq 2$, we introduce a set of triplets

$$\mathcal{I}_\ell = \mathcal{I}_\ell(X_r) := \{(a, m, u) \mid a \in I; m = 1, \dots, t_a\ell - 1; u \in \frac{1}{t}\mathbb{Z}\}. \tag{2.3}$$

Definition 2.1 ([KNS]). Fix an integer $\ell \geq 2$. The *level ℓ restricted T-system $\mathbb{T}_\ell(X_r)$ of type X_r (with the unit boundary condition)* is the following system of relations for a family of variables $T_\ell = \{T^{(a)}_m(u) \mid (a, m, u) \in \mathcal{I}_\ell\}$, where $T^{(0)}_m(u) = T^{(a)}_0(u) = 1$, and furthermore, $T^{(a)}_{t_a\ell}(u) = 1$ (the *unit boundary condition*) if they occur in the right hand sides in the relations:

(Here and throughout the paper, $2m$ (resp. $2m+1$) in the left hand sides, for example, represents elements $2, 4, \dots$ (resp. $1, 3, \dots$).)

For $X_r = C_r$,

$$\begin{aligned}
T^{(a)}_m\left(u-\tfrac{1}{2}\right)T^{(a)}_m\left(u+\tfrac{1}{2}\right) &= T^{(a)}_{m-1}(u)T^{(a)}_{m+1}(u) + T^{(a-1)}_m(u)T^{(a+1)}_m(u)\\
&\qquad\qquad\qquad\qquad (1 \le a \le r-2),\\
T^{(r-1)}_{2m}\left(u-\tfrac{1}{2}\right)T^{(r-1)}_{2m}\left(u+\tfrac{1}{2}\right) &= T^{(r-1)}_{2m-1}(u)T^{(r-1)}_{2m+1}(u)\\
&\quad + T^{(r-2)}_{2m}(u)T^{(r)}_m\left(u-\tfrac{1}{2}\right)T^{(r)}_m\left(u+\tfrac{1}{2}\right),\\
T^{(r-1)}_{2m+1}\left(u-\tfrac{1}{2}\right)T^{(r-1)}_{2m+1}\left(u+\tfrac{1}{2}\right) &= T^{(r-1)}_{2m}(u)T^{(r-1)}_{2m+2}(u)\\
&\quad + T^{(r-2)}_{2m+1}(u)T^{(r)}_m(u)T^{(r)}_{m+1}(u),\\
T^{(r)}_m(u-1)T^{(r)}_m(u+1) &= T^{(r)}_{m-1}(u)T^{(r)}_{m+1}(u) + T^{(r-1)}_{2m}(u).
\end{aligned}\tag{2.4}$$

For $X_r = F_4$,

$$\begin{aligned}
T^{(1)}_m(u-1)T^{(1)}_m(u+1) &= T^{(1)}_{m-1}(u)T^{(1)}_{m+1}(u) + T^{(2)}_m(u),\\
T^{(2)}_m(u-1)T^{(2)}_m(u+1) &= T^{(2)}_{m-1}(u)T^{(2)}_{m+1}(u) + T^{(1)}_m(u)T^{(3)}_{2m}(u),\\
T^{(3)}_{2m}\left(u-\tfrac{1}{2}\right)T^{(3)}_{2m}\left(u+\tfrac{1}{2}\right) &= T^{(3)}_{2m-1}(u)T^{(3)}_{2m+1}(u)\\
&\quad + T^{(2)}_m\left(u-\tfrac{1}{2}\right)T^{(2)}_m\left(u+\tfrac{1}{2}\right)T^{(4)}_{2m}(u),\\
T^{(3)}_{2m+1}\left(u-\tfrac{1}{2}\right)T^{(3)}_{2m+1}\left(u+\tfrac{1}{2}\right) &= T^{(3)}_{2m}(u)T^{(3)}_{2m+2}(u) + T^{(2)}_m(u)T^{(2)}_{m+1}(u)T^{(4)}_{2m+1}(u),\\
T^{(4)}_m\left(u-\tfrac{1}{2}\right)T^{(4)}_m\left(u+\tfrac{1}{2}\right) &= T^{(4)}_{m-1}(u)T^{(4)}_{m+1}(u) + T^{(3)}_m(u).
\end{aligned}\tag{2.5}$$

For $X_r = G_2$,

$$\begin{aligned}
T^{(1)}_m(u-1)T^{(1)}_m(u+1) &= T^{(1)}_{m-1}(u)T^{(1)}_{m+1}(u) + T^{(2)}_{3m}(u),\\
T^{(2)}_{3m}\left(u-\tfrac{1}{3}\right)T^{(2)}_{3m}\left(u+\tfrac{1}{3}\right) &= T^{(2)}_{3m-1}(u)T^{(2)}_{3m+1}(u)\\
&\quad + T^{(1)}_m\left(u-\tfrac{2}{3}\right)T^{(1)}_m(u)T^{(1)}_m\left(u+\tfrac{2}{3}\right),\\
T^{(2)}_{3m+1}\left(u-\tfrac{1}{3}\right)T^{(2)}_{3m+1}\left(u+\tfrac{1}{3}\right) &= T^{(2)}_{3m}(u)T^{(2)}_{3m+2}(u)\\
&\quad + T^{(1)}_m\left(u-\tfrac{1}{3}\right)T^{(1)}_m\left(u+\tfrac{1}{3}\right)T^{(1)}_{m+1}(u),\\
T^{(2)}_{3m+2}\left(u-\tfrac{1}{3}\right)T^{(2)}_{3m+2}\left(u+\tfrac{1}{3}\right) &= T^{(2)}_{3m+1}(u)T^{(2)}_{3m+3}(u)\\
&\quad + T^{(1)}_m(u)T^{(1)}_{m+1}\left(u-\tfrac{1}{3}\right)T^{(1)}_{m+1}\left(u+\tfrac{1}{3}\right).
\end{aligned}\tag{2.6}$$

Definition 2.2 ([KN])**.** Fix an integer $\ell \ge 2$. The *level ℓ restricted Y-system $\mathbb{Y}_\ell(X_r)$ of type X_r* is the following system of relations for a family of variables $Y_\ell = \{Y^{(a)}_m(u) \mid (a,m,u) \in \mathcal{I}_\ell\}$, where $Y^{(0)}_m(u) = Y^{(a)}_0(u)^{-1} = Y^{(a)}_{t_a\ell}(u)^{-1} = 0$ if they occur in the right hand sides in the relations:

For $X_r = C_r$,

(2.7)
$$\begin{aligned}
Y^{(a)}_m\left(u-\tfrac{1}{2}\right)Y^{(a)}_m\left(u+\tfrac{1}{2}\right) &= \frac{(1+Y^{(a-1)}_m(u))(1+Y^{(a+1)}_m(u))}{(1+Y^{(a)}_{m-1}(u)^{-1})(1+Y^{(a)}_{m+1}(u)^{-1})}\\
&\qquad\qquad\qquad\qquad\qquad (1\le a\le r-2),\\
Y^{(r-1)}_{2m}\left(u-\tfrac{1}{2}\right)Y^{(r-1)}_{2m}\left(u+\tfrac{1}{2}\right) &= \frac{(1+Y^{(r-2)}_{2m}(u))(1+Y^{(r)}_m(u))}{(1+Y^{(r-1)}_{2m-1}(u)^{-1})(1+Y^{(r-1)}_{2m+1}(u)^{-1})},\\
Y^{(r-1)}_{2m+1}\left(u-\tfrac{1}{2}\right)Y^{(r-1)}_{2m+1}\left(u+\tfrac{1}{2}\right) &= \frac{1+Y^{(r-2)}_{2m+1}(u)}{(1+Y^{(r-1)}_{2m}(u)^{-1})(1+Y^{(r-1)}_{2m+2}(u)^{-1})},\\
Y^{(r)}_m(u-1)Y^{(r)}_m(u+1) &= \frac{\begin{array}{c}(1+Y^{(r-1)}_{2m+1}(u))(1+Y^{(r-1)}_{2m-1}(u))\\ \times(1+Y^{(r-1)}_{2m}\left(u-\tfrac{1}{2}\right))(1+Y^{(r-1)}_{2m}\left(u+\tfrac{1}{2}\right))\end{array}}{(1+Y^{(r)}_{m-1}(u)^{-1})(1+Y^{(r)}_{m+1}(u)^{-1})}.
\end{aligned}$$

For $X_r = F_4$,

(2.8)
$$\begin{aligned}
Y^{(1)}_m(u-1)Y^{(1)}_m(u+1) &= \frac{1+Y^{(2)}_m(u)}{(1+Y^{(1)}_{m-1}(u)^{-1})(1+Y^{(1)}_{m+1}(u)^{-1})},\\
Y^{(2)}_m(u-1)Y^{(2)}_m(u+1) &= \frac{\begin{array}{c}(1+Y^{(1)}_m(u))(1+Y^{(3)}_{2m-1}(u))(1+Y^{(3)}_{2m+1}(u))\\ \times(1+Y^{(3)}_{2m}\left(u-\tfrac{1}{2}\right))(1+Y^{(3)}_{2m}\left(u+\tfrac{1}{2}\right))\end{array}}{(1+Y^{(2)}_{m-1}(u)^{-1})(1+Y^{(2)}_{m+1}(u)^{-1})},\\
Y^{(3)}_{2m}\left(u-\tfrac{1}{2}\right)Y^{(3)}_{2m}\left(u+\tfrac{1}{2}\right) &= \frac{(1+Y^{(2)}_m(u))(1+Y^{(4)}_{2m}(u))}{(1+Y^{(3)}_{2m-1}(u)^{-1})(1+Y^{(3)}_{2m+1}(u)^{-1})},\\
Y^{(3)}_{2m+1}\left(u-\tfrac{1}{2}\right)Y^{(3)}_{2m+1}\left(u+\tfrac{1}{2}\right) &= \frac{1+Y^{(4)}_{2m+1}(u)}{(1+Y^{(3)}_{2m}(u)^{-1})(1+Y^{(3)}_{2m+2}(u)^{-1})},\\
Y^{(4)}_m\left(u-\tfrac{1}{2}\right)Y^{(4)}_m\left(u+\tfrac{1}{2}\right) &= \frac{1+Y^{(3)}_m(u)}{(1+Y^{(4)}_{m-1}(u)^{-1})(1+Y^{(4)}_{m+1}(u)^{-1})}.
\end{aligned}$$

For $X_r = G_2$,

(2.9)
$$
\begin{aligned}
Y^{(1)}_m(u-1)Y^{(1)}_m(u+1) &= \frac{\begin{array}{l}(1+Y^{(2)}_{3m-2}(u))(1+Y^{(2)}_{3m+2}(u))\\ \times(1+Y^{(2)}_{3m-1}\left(u-\frac{1}{3}\right))(1+Y^{(2)}_{3m-1}\left(u+\frac{1}{3}\right))\\ \times(1+Y^{(2)}_{3m+1}\left(u-\frac{1}{3}\right))(1+Y^{(2)}_{3m+1}\left(u+\frac{1}{3}\right))\\ \times(1+Y^{(2)}_{3m}\left(u-\frac{2}{3}\right))(1+Y^{(2)}_{3m}\left(u+\frac{2}{3}\right))\\ \times(1+Y^{(2)}_{3m}(u))\end{array}}{(1+Y^{(1)}_{m-1}(u)^{-1})(1+Y^{(1)}_{m+1}(u)^{-1})},\\
Y^{(2)}_{3m}\left(u-\tfrac{1}{3}\right)Y^{(2)}_{3m}\left(u+\tfrac{1}{3}\right) &= \frac{1+Y^{(1)}_m(u)}{(1+Y^{(2)}_{3m-1}(u)^{-1})(1+Y^{(2)}_{3m+1}(u)^{-1})},\\
Y^{(2)}_{3m+1}\left(u-\tfrac{1}{3}\right)Y^{(2)}_{3m+1}\left(u+\tfrac{1}{3}\right) &= \frac{1}{(1+Y^{(2)}_{3m}(u)^{-1})(1+Y^{(2)}_{3m+2}(u)^{-1})},\\
Y^{(2)}_{3m+2}\left(u-\tfrac{1}{3}\right)Y^{(2)}_{3m+2}\left(u+\tfrac{1}{3}\right) &= \frac{1}{(1+Y^{(2)}_{3m+1}(u)^{-1})(1+Y^{(2)}_{3m+3}(u)^{-1})}.
\end{aligned}
$$

Let us write (2.4)–(2.6) in a unified manner

(2.10)
$$
T^{(a)}_m\left(u-\frac{1}{t_a}\right)T^{(a)}_m\left(u+\frac{1}{t_a}\right) = T^{(a)}_{m-1}(u)T^{(a)}_{m+1}(u) + \prod_{(b,k,v)\in\mathcal{I}_\ell} T^{(b)}_k(v)^{G(b,k,v;a,m,u)}.
$$

Define the transposition ${}^tG(b,k,v;a,m,u) = G(a,m,u;b,k,v)$. Then, (2.7)–(2.9) are written as

$$
Y^{(a)}_m\left(u-\frac{1}{t_a}\right)Y^{(a)}_m\left(u+\frac{1}{t_a}\right) = \frac{\displaystyle\prod_{(b,k,v)\in\mathcal{I}_\ell}(1+Y^{(b)}_k(v))^{{}^tG(b,k,v;a,m,u)}}{(1+Y^{(a)}_{m-1}(u)^{-1})(1+Y^{(a)}_{m+1}(u)^{-1})}. \tag{2.11}
$$

2.2. **Periodicities.**

Definition 2.3. Let $\mathcal{T}_\ell(X_r)$ be the commutative ring over $\mathbb{Z}$ with identity element, with generators $T^{(a)}_m(u)^{\pm1}$ ($(a,m,u)\in\mathcal{I}_\ell$) and relations $\mathbb{T}_\ell(X_r)$ together with $T^{(a)}_m(u)T^{(a)}_m(u)^{-1}=1$. Let $\mathcal{T}^\circ_\ell(X_r)$ be the subring of $\mathcal{T}_\ell(X_r)$ generated by $T^{(a)}_m(u)$ ($(a,m,u)\in\mathcal{I}_\ell$).

Definition 2.4. Let $\mathcal{Y}_\ell(X_r)$ be the semifield with generators $Y^{(a)}_m(u)$ ($(a,m,u)\in\mathcal{I}_\ell$ and relations $\mathbb{Y}_\ell(X_r)$. Let $\mathcal{Y}^\circ_\ell(X_r)$ be the multiplicative subgroup of $\mathcal{Y}_\ell(X_r)$ generated by $Y^{(a)}_m(u)$, $1+Y^{(a)}_m(u)$ ($(a,m,u)\in\mathcal{I}_\ell$). (Here we use the symbol $+$ instead of $\oplus$ for simplicity.)

The first main result of the paper is the periodicities of the T and Y-systems.

Theorem 2.5 (Conjectured by [IIKNS]). *The following relations hold in $\mathcal{T}^\circ_\ell(X_r)$.*

(i) Half periodicity: $T^{(a)}_m(u+h^\vee+\ell) = T^{(a)}_{t_a\ell-m}(u)$.

(ii) Full periodicity: $T^{(a)}_m(u+2(h^\vee+\ell)) = T^{(a)}_m(u)$.

Theorem 2.6 (Conjectured by [KNS]). *The following relations hold in $\mathcal{Y}^\circ_\ell(X_r)$.*

(i) Half periodicity: $Y^{(a)}_m(u+h^\vee+\ell) = Y^{(a)}_{t_a\ell-m}(u)$.

(ii) Full periodicity: $Y^{(a)}_m(u+2(h^\vee+\ell)) = Y^{(a)}_m(u)$.

2.3. **Dilogarithm identities.** Let $L(x)$ be the *Rogers dilogarithm function*

$$L(x) = -\frac{1}{2}\int_0^x \left\{ \frac{\log(1-y)}{y} + \frac{\log y}{1-y} \right\} dy \quad (0 \le x \le 1). \tag{2.12}$$

We introduce the *constant version* of the Y-system.

Definition 2.7. Fix an integer $\ell \ge 2$. The *level ℓ restricted constant Y-system $\mathbb{Y}^{\mathrm{c}}_\ell(X_r)$ of type X_r* is the following system of relations for a family of variables $Y^{\mathrm{c}}_\ell = \{Y^{(a)}_m \mid a \in I; m = 1, \dots, t_a\ell - 1\}$, where $Y^{(0)}_m = Y^{(a)}_0{}^{-1} = Y^{(a)}_{t_a\ell}{}^{-1} = 0$ if they occur in the right hand sides in the relations:

For $X_r = C_r$,

$$\begin{aligned}
(Y^{(a)}_m)^2 &= \frac{(1+Y^{(a-1)}_m)(1+Y^{(a+1)}_m)}{(1+Y^{(a)}_{m-1}{}^{-1})(1+Y^{(a)}_{m+1}{}^{-1})} \quad (1 \le a \le r-2),\\
(Y^{(r-1)}_{2m})^2 &= \frac{(1+Y^{(r-2)}_{2m})(1+Y^{(r)}_m)}{(1+Y^{(r-1)}_{2m-1}{}^{-1})(1+Y^{(r-1)}_{2m+1}{}^{-1})},\\
(Y^{(r-1)}_{2m+1})^2 &= \frac{1+Y^{(r-2)}_{2m+1}}{(1+Y^{(r-1)}_{2m}{}^{-1})(1+Y^{(r-1)}_{2m+2}{}^{-1})},\\
(Y^{(r)}_m)^2 &= \frac{(1+Y^{(r-1)}_{2m-1})(1+Y^{(r-1)}_{2m})^2(1+Y^{(r-1)}_{2m+1})}{(1+Y^{(r)}_{m-1}{}^{-1})(1+Y^{(r)}_{m+1}{}^{-1})}.
\end{aligned} \tag{2.13}$$

For $X_r = F_4$,

$$\begin{aligned}
(Y^{(1)}_m)^2 &= \frac{1+Y^{(2)}_m}{(1+Y^{(1)}_{m-1}{}^{-1})(1+Y^{(1)}_{m+1}{}^{-1})},\\
(Y^{(2)}_m)^2 &= \frac{(1+Y^{(1)}_m)(1+Y^{(3)}_{2m-1})(1+Y^{(3)}_{2m})^2(1+Y^{(3)}_{2m+1})}{(1+Y^{(2)}_{m-1}(u)^{-1})(1+Y^{(2)}_{m+1}(u)^{-1})},\\
(Y^{(3)}_{2m})^2 &= \frac{(1+Y^{(2)}_m)(1+Y^{(4)}_{2m})}{(1+Y^{(3)}_{2m-1}{}^{-1})(1+Y^{(3)}_{2m+1}{}^{-1})},\\
(Y^{(3)}_{2m+1})^2 &= \frac{1+Y^{(4)}_{2m+1}}{(1+Y^{(3)}_{2m}{}^{-1})(1+Y^{(3)}_{2m+2}{}^{-1})},\\
(Y^{(4)}_m)^2 &= \frac{1+Y^{(3)}_m}{(1+Y^{(4)}_{m-1}{}^{-1})(1+Y^{(4)}_{m+1}{}^{-1})}.
\end{aligned} \tag{2.14}$$

For $X_r = G_2$,

(2.15)
$$\begin{aligned}
(Y^{(1)}_m)^2 &= \frac{(1+Y^{(2)}_{3m-2})(1+Y^{(2)}_{3m-1})^2(1+Y^{(2)}_{3m})^3(1+Y^{(2)}_{3m+1})^2(1+Y^{(2)}_{3m+2})}{(1+Y^{(1)}_{m-1}{}^{-1})(1+Y^{(1)}_{m+1}{}^{-1})},\\
(Y^{(2)}_{3m})^2 &= \frac{1+Y^{(1)}_m}{(1+Y^{(2)}_{3m-1}{}^{-1})(1+Y^{(2)}_{3m+1}{}^{-1})},\\
(Y^{(2)}_{3m+1})^2 &= \frac{1}{(1+Y^{(2)}_{3m}{}^{-1})(1+Y^{(2)}_{3m+2}{}^{-1})},\\
(Y^{(2)}_{3m+2})^2 &= \frac{1}{(1+Y^{(2)}_{3m+1}{}^{-1})(1+Y^{(2)}_{3m+3}{}^{-1})}.
\end{aligned}$$

Proposition 2.8. *There exists a unique positive real solution of $\mathbb{Y}^{\mathrm{c}}_\ell(X_r)$.*

Proof. The same proof of [IIKKN, Proposition 1.8] is applicable. □

The second main result of the paper is the dilogarithm identities conjectured by Kirillov [Ki, Eq. (7)], properly corrected by Kuniba [Ku, Eqs. (A.1a), (A.1c)],

Theorem 2.9 (Dilogarithm identities)**.** *Suppose that a family of positive real numbers $\{Y^{(a)}_m \mid a \in I; m = 1,\dots,t_a\ell-1\}$ satisfies $\mathbb{Y}^{\mathrm{c}}_\ell(X_r)$. Then, we have the identity*

$$\frac{6}{\pi^2}\sum_{a\in I}\sum_{m=1}^{t_a\ell-1} L\left(\frac{Y^{(a)}_m}{1+Y^{(a)}_m}\right) = \frac{\ell \dim \mathfrak{g}}{h^\vee+\ell} - r, \tag{2.16}$$

where $\mathfrak{g}$ is the simple Lie algebra of type X_r.

The right hand side of (2.16) is equal to the number

$$\frac{r(\ell h - h^\vee)}{h^\vee+\ell}. \tag{2.17}$$

In fact, we prove a functional generalization of Theorem 2.9.

Theorem 2.10 (Functional dilogarithm identities)**.** *Suppose that a family of positive real numbers $\{Y^{(a)}_m(u) \mid (a,m,u)\in\mathcal{I}_\ell\}$ satisfies $\mathbb{Y}_\ell(X_r)$. Then, we have the identities*

$$\begin{aligned}
\frac{6}{\pi^2}\sum_{\substack{(a,m,u)\in\mathcal{I}_\ell\\ 0\le u<2(h^\vee+\ell)}} L\left(\frac{Y^{(a)}_m(u)}{1+Y^{(a)}_m(u)}\right) &= 2tr(\ell h - h^\vee)\\
&= \begin{cases} 4r(2r\ell-r-1) & C_r\\ 48(4\ell-3) & F_4\\ 24(3\ell-2) & G_2,\end{cases}
\end{aligned} \tag{2.18}$$

$$\frac{6}{\pi^2}\sum_{\substack{(a,m,u)\in\mathcal{I}_\ell\\ 0\le u<2(h^\vee+\ell)}} L\left(\frac{1}{1+Y^{(a)}_m(u)}\right) = \begin{cases} 4\ell(2r\ell-\ell-1) & C_r\\ 8\ell(3\ell+1) & F_4\\ 12\ell(2\ell+1) & G_2.\end{cases} \tag{2.19}$$



FIGURE 1. The quiver $Q_\ell(C_r)$ for even ℓ (upper) and for odd ℓ (lower), where we identify the rightmost column in the left quiver with the middle column in the right quiver.

The two identities (2.18) and (2.19) are equivalent to each other, since the sum of the right hand sides is equal to $2t(h^\vee+\ell)((\sum_{a\in I} t_a)\ell - r)$, which is the total number of $(a,m,u)\in I_\ell$ in the region $0\leq u<2(h^\vee+\ell)$.

It is clear that Theorem 2.9 follows form Theorem 2.10.

3. TYPE C_r

The C_r case is quite parallel to the B_r case. For the reader's convenience, we repeat most of the basic definitions and results in [IIKKN]. Most propositions are proved in a parallel manner to the B_r case, so that proofs are omitted. The properties of the tropical Y-system at level 2 (Proposition 3.10) are crucial and specific to C_r. Since its derivation is a little more complicated than the B_r case, the outline of the proof is provided.

3.1. **Parity decompositions of T and Y-systems.**

For a triplet $(a,m,u)\in\mathcal{I}_\ell$, we set the 'parity conditions' $\mathbf{P}_+$ and $\mathbf{P}_-$ by

$$\mathbf{P}_+:\ r+a+m+2u \text{ is odd if } a\neq r;\ 2u \text{ is even if } a=r, \tag{3.1}$$

$$\mathbf{P}_-:\ r+a+m+2u \text{ is even if } a\neq r;\ 2u \text{ is odd if } a=r. \tag{3.2}$$

We write, for example, $(a,m,u):\mathbf{P}_+$ if (a,m,u) satisfies $\mathbf{P}_+$. We have $\mathcal{I}_\ell=\mathcal{I}_{\ell+}\sqcup\mathcal{I}_{\ell-}$, where $\mathcal{I}_{\ell\varepsilon}$ is the set of all $(a,m,u):\mathbf{P}_\varepsilon$.

Define $\mathcal{T}^\circ_\ell(C_r)_\varepsilon$ ($\varepsilon = \pm$) to be the subring of $\mathcal{T}^\circ_\ell(C_r)$ generated by $T^{(a)}_m(u)$ ($(a,m,u) \in \mathcal{I}_{\ell\varepsilon}$). Then, we have $\mathcal{T}^\circ_\ell(C_r)_+ \simeq \mathcal{T}^\circ_\ell(C_r)_-$ by $T^{(a)}_m(u) \mapsto T^{(a)}_m(u+\frac{1}{2})$ and

$$\mathcal{T}^\circ_\ell(C_r) \simeq \mathcal{T}^\circ_\ell(C_r)_+ \otimes_{\mathbb{Z}} \mathcal{T}^\circ_\ell(C_r)_-. \tag{3.3}$$

For a triplet $(a,m,u) \in \mathcal{I}_\ell$, we set another 'parity conditions' $\mathbf{P}'_+$ and $\mathbf{P}'_-$ by

$$\mathbf{P}'_+ : \ r+a+m+2u \text{ is even if } a \neq r;\ 2u \text{ is even if } a = r, \tag{3.4}$$

$$\mathbf{P}'_- : \ r+a+m+2u \text{ is odd if } a \neq r;\ 2u \text{ is odd if } a = r. \tag{3.5}$$

We have $\mathcal{I}_\ell = \mathcal{I}'_{\ell+} \sqcup \mathcal{I}'_{\ell-}$, where $\mathcal{I}'_{\ell\varepsilon}$ is the set of all $(a,m,u) : \mathbf{P}'_\varepsilon$. We also have

$$(a,m,u) : \mathbf{P}'_+ \iff (a,m,u \pm \tfrac{1}{t_a}) : \mathbf{P}_+. \tag{3.6}$$

Define $\mathcal{Y}^\circ_\ell(C_r)_\varepsilon$ ($\varepsilon = \pm$) to be the subgroup of $\mathcal{Y}^\circ_\ell(C_r)$ generated by $Y^{(a)}_m(u)$, $1+Y^{(a)}_m(u)$ ($(a,m,u) \in \mathcal{I}'_{\ell\varepsilon}$). Then, we have $\mathcal{Y}^\circ_\ell(C_r)_+ \simeq \mathcal{Y}^\circ_\ell(C_r)_-$ by $Y^{(a)}_m(u) \mapsto Y^{(a)}_m(u+\frac{1}{2})$, $1+Y^{(a)}_m(u) \mapsto 1+Y^{(a)}_m(u+\frac{1}{2})$, and

$$\mathcal{Y}^\circ_\ell(C_r) \simeq \mathcal{Y}^\circ_\ell(C_r)_+ \times \mathcal{Y}^\circ_\ell(C_r)_-. \tag{3.7}$$

3.2. Quiver $Q_\ell(C_r)$.

With type C_r and $\ell \geq 2$ we associate the quiver $Q_\ell(C_r)$ by Figure 1, where the rightmost column in the left quiver and the middle column in the right quiver are identified. Also, we assign the empty or filled circle $\circ/\bullet$ and the sign $+/-$ to each vertex.

Let us choose the index set $\mathbf{I}$ of the vertices of $Q_\ell(C_r)$ so that $\mathbf{i} = (i,i') \in \mathbf{I}$ represents the vertex at the i'th row (from the bottom) of the ith column (from the left) in the left quiver for $i = 1,\dots,r-1$, the one of the right column in the right quiver for $i = r$, and the one of the left column in the right quiver for $i = r+1$. Thus, $i = 1,\dots,r+1$, and $i' = 1,\dots,\ell-1$ if $i \neq r, r+1$ and $i' = 1,\dots,2\ell-1$ if $i = r, r+1$. We use a natural notation $\mathbf{I}^\circ$ (resp. $\mathbf{I}^\circ_+$) for the set of the vertices $\mathbf{i}$ with property $\circ$ (resp. $\circ$ and $+$), and so on. We have $\mathbf{I} = \mathbf{I}^\circ \sqcup \mathbf{I}^\bullet = \mathbf{I}^\circ_+ \sqcup \mathbf{I}^\circ_- \sqcup \mathbf{I}^\bullet_+ \sqcup \mathbf{I}^\bullet_-$.

We define composite mutations,

$$\mu^\circ_+ = \prod_{\mathbf{i}\in\mathbf{I}^\circ_+} \mu_{\mathbf{i}}, \quad \mu^\circ_- = \prod_{\mathbf{i}\in\mathbf{I}^\circ_-} \mu_{\mathbf{i}}, \quad \mu^\bullet_+ = \prod_{\mathbf{i}\in\mathbf{I}^\bullet_+} \mu_{\mathbf{i}}, \quad \mu^\bullet_- = \prod_{\mathbf{i}\in\mathbf{I}^\bullet_-} \mu_{\mathbf{i}}. \tag{3.8}$$

Note that they do not depend on the order of the product.

Let $\boldsymbol{r}$ be the involution acting on $\mathbf{I}$ by the left-right reflection of the right quiver. Let $\boldsymbol{\omega}$ be the involution acting on $\mathbf{I}$ defined by, for even r, the up-down reflection of the left quiver and the 180° rotation of the right quiver; and for odd r, the up-down reflection of the left and right quivers. Let $\boldsymbol{r}(Q_\ell(C_r))$ and $\boldsymbol{\omega}(Q_\ell(C_r))$ denote the quivers induced from $Q_\ell(C_r)$ by $\boldsymbol{r}$ and $\boldsymbol{\omega}$, respectively. For example, if there is an arrow $\mathbf{i} \to \mathbf{j}$ in $Q_\ell(C_r)$, then, there is an arrow $\boldsymbol{r}(\mathbf{i}) \to \boldsymbol{r}(\mathbf{j})$ in $\boldsymbol{r}(Q_\ell(C_r))$. For a quiver Q, Q^{op} denotes the opposite quiver.

Lemma 3.1. *Let $Q = Q_\ell(C_r)$.*

(i) We have a periodic sequence of mutations of quivers

$$Q \overset{\mu^\bullet_+\mu^\circ_+}{\longleftrightarrow} Q^{\mathrm{op}} \overset{\mu^\bullet_-}{\longleftrightarrow} \boldsymbol{r}(Q) \overset{\mu^\bullet_+\mu^\circ_-}{\longleftrightarrow} \boldsymbol{r}(Q)^{\mathrm{op}} \overset{\mu^\bullet_-}{\longleftrightarrow} Q. \tag{3.9}$$

(ii) $\boldsymbol{\omega}(Q) = Q$ if $h^\vee + \ell$ is even, and $\boldsymbol{\omega}(Q) = \boldsymbol{r}(Q)$ if $h^\vee + \ell$ is odd.


$x_7^{(r-1)}(-\frac{1}{2})$
$x_6^{(r-2)}(-\frac{1}{2})$
$x_3^{(r)}(-1)$
$x_5^{(r-1)}(-\frac{1}{2})$
$x(0)$
$x_4^{(r-2)}(-\frac{1}{2})$
$x_2^{(r)}(-1)$
$x_3^{(r-1)}(-\frac{1}{2})$
$x_2^{(r-2)}(-\frac{1}{2})$
$x_1^{(r)}(-1)$
$x_1^{(r-1)}(-\frac{1}{2})$
$\mu_+^\bullet \mu_+^\circ \updownarrow$
$x_7^{(r-2)}(0)$
$x_6^{(r-1)}(0)$
$x_5^{(r-2)}(0)$
$x(\frac{1}{2})$
$x_4^{(r-1)}(0)$
$x_3^{(r-2)}(0)$
$x_2^{(r-1)}(0)$
$x_1^{(r-2)}(0)$
$\mu_-^\bullet \updownarrow$

FIGURE 2. (Continues to Figure 3) Label of cluster variables $x_{\mathbf{i}}(u)$ by $\mathcal{I}_{\ell+}$ for C_r, $\ell = 4$. The variables framed by solid/dashed lines satisfy the condition $\mathbf{p}_+/\mathbf{p}_-$, respectively. The middle column in the right quiver (marked by $\diamond$) is identified with the rightmost column in the left quiver.

3.3. **Cluster algebra and alternative labels.** It is standard to identify a quiver Q with no loop and no 2-cycle and a skew-symmetric matrix B. We use the convention for the direction of arrows as

$$i \longrightarrow j \quad \Longleftrightarrow \quad B_{ij} = 1. \tag{3.10}$$

(In this paper we only encounter the situation where $B_{ij} = -1, 0, 1$.) Let $B_\ell(C_r)$ be the corresponding skew-symmetric matrix to the quiver $Q_\ell(C_r)$. In the rest of the section, we set the matrix $B = (B_{\mathbf{ij}})_{\mathbf{i},\mathbf{j}\in\mathbf{I}} = B_\ell(C_r)$ unless otherwise mentioned.

Let $\mathcal{A}(B, x, y)$ be the *cluster algebra with coefficients in the universal semifield* $\mathbb{Q}_{\mathrm{sf}}(y)$, where (B, x, y) is the initial seed [FZ2]. See also [IIKKN, Section 2.1] for the conventions and notations on cluster algebras we employ. (Here we use the



FIGURE 3. (Continues from Figure 2).

symbol $+$ instead of $\oplus$ in $\mathbb{Q}_{\mathrm{sf}}(y)$, since it is the ordinary addition of subtraction-free expressions of rational functions of y.)

Definition 3.2. The *coefficient group* $\mathcal{G}(B,y)$ *associated with* $\mathcal{A}(B,x,y)$ is the multiplicative subgroup of the semifield $\mathbb{Q}_{\mathrm{sf}}(y)$ generated by all the coefficients $y'_{\mathbf{i}}$ of $\mathcal{A}(B,x,y)$ together with $1+y'_{\mathbf{i}}$.

In view of Lemma 3.1 we set $x(0)=x$, $y(0)=y$ and define clusters $x(u)=(x_{\mathbf{i}}(u))_{\mathbf{i}\in\mathbf{I}}$ ($u\in\frac12\mathbb{Z}$) and coefficient tuples $y(u)=(y_{\mathbf{i}}(u))_{\mathbf{i}\in\mathbf{I}}$ ($u\in\frac12\mathbb{Z}$) by the sequence of mutations

$$\begin{aligned}(3.11)\qquad &\cdots\overset{\mu_-^\bullet}{\longleftrightarrow}(B,x(0),y(0))\overset{\mu_+^\bullet\mu_+^\circ}{\longleftrightarrow}(-B,x(\tfrac12),y(\tfrac12))\\ &\overset{\mu_-^\bullet}{\longleftrightarrow}(\boldsymbol{r}(B),x(1),y(1))\overset{\mu_+^\bullet\mu_-^\circ}{\longleftrightarrow}(-\boldsymbol{r}(B),x(\tfrac32),y(\tfrac32))\overset{\mu_-^\bullet}{\longleftrightarrow}\cdots,\end{aligned}$$

where $\boldsymbol{r}(B)=B'$ is defined by $B'_{\mathbf{ij}}=B_{\boldsymbol{r}(\mathbf{i})\boldsymbol{r}(\mathbf{j})}$.

For a pair $(\mathbf{i}, u) \in \mathbf{I} \times \frac{1}{2}\mathbb{Z}$, we set the parity condition $\mathbf{p}_+$ and $\mathbf{p}_-$ by

$$\mathbf{p}_+ : \begin{cases} \mathbf{i} \in \mathbf{I}^\circ_+ \sqcup \mathbf{I}^\bullet_+ & u \equiv 0 \\ \mathbf{i} \in \mathbf{I}^\bullet_- & u \equiv \frac{1}{2}, \frac{3}{2} \\ \mathbf{i} \in \mathbf{I}^\circ_- \sqcup \mathbf{I}^\bullet_+ & u \equiv 1, \end{cases} \qquad \mathbf{p}_- : \begin{cases} \mathbf{i} \in \mathbf{I}^\circ_+ \sqcup \mathbf{I}^\bullet_+ & u \equiv \frac{1}{2} \\ \mathbf{i} \in \mathbf{I}^\bullet_- & u \equiv 0, 1 \\ \mathbf{i} \in \mathbf{I}^\circ_- \sqcup \mathbf{I}^\bullet_+ & u \equiv \frac{3}{2}, \end{cases} \tag{3.12}$$

where $\equiv$ is modulo $2\mathbb{Z}$. We have

$$(\mathbf{i}, u) : \mathbf{p}_+ \iff (\mathbf{i}, u + \frac{1}{2}) : \mathbf{p}_-. \tag{3.13}$$

Each $(\mathbf{i}, u) : \mathbf{p}_+$ is a mutation point of (3.11) in the forward direction of u, and each $(\mathbf{i}, u) : \mathbf{p}_-$ is so in the backward direction of u. Notice that there are also some $(\mathbf{i}, u)$ which do not satisfy $\mathbf{p}_+$ nor $\mathbf{p}_-$, and are not mutation points of (3.11); explicitly, they are $(\mathbf{i}, u)$ with $\mathbf{i} \in \mathbf{I}^\circ_+$, $u \equiv 1, \frac{3}{2}$ mod $2\mathbb{Z}$, or with $\mathbf{i} \in \mathbf{I}^\circ_-$, $u \equiv 0, \frac{1}{2}$ mod $2\mathbb{Z}$.

There is a correspondence between the parity condition $\mathbf{p}_\pm$ here and $\mathbf{P}_\pm$, $\mathbf{P}'_\pm$ in (3.1) and (3.4).

Lemma 3.3. *Below $\equiv$ means the equivalence modulo $2\mathbb{Z}$.*

(i) The map

$$\begin{array}{rccl} g: & \mathcal{I}_{\ell+} & \to & \{(\mathbf{i}, u) : \mathbf{p}_+\} \\ & (a, m, u - \frac{1}{t_a}) & \mapsto & \begin{cases} ((a, m), u) & a \neq r \\ ((r+1, m), u) & a = r;\ m + u \equiv 0 \\ ((r, m), u) & a = r;\ m + u \equiv 1 \end{cases} \end{array} \tag{3.14}$$

is a bijection.

(ii) The map

$$\begin{array}{rccl} g': & \mathcal{I}'_{\ell+} & \to & \{(\mathbf{i}, u) : \mathbf{p}_+ \text{ or } \mathbf{p}_-\} \\ & (a, m, u) & \mapsto & \begin{cases} ((a, m), u) & a \neq r \\ ((r+1, m), u) & a = r;\ m + u \equiv 0 \\ ((r, m), u) & a = r;\ m + u \equiv 1 \end{cases} \end{array} \tag{3.15}$$

is a bijection.

We introduce alternative labels $x_{\mathbf{i}}(u) = x^{(a)}_m(u - 1/t_a)$ $((a, m, u - 1/t_a) \in \mathcal{I}_{\ell+})$ for $(\mathbf{i}, u) = g((a, m, u - 1/t_a))$ and $y_{\mathbf{i}}(u) = y^{(a)}_m(u)$ $((a, m, u) \in \mathcal{I}'_{\ell+})$ for $(\mathbf{i}, u) = g'((a, m, u))$, respectively. See Figures 2–3.

3.4. **T-system and cluster algebra.**

The result in this subsection is completely parallel to the B_r case [IIKKN].

Let $\mathcal{A}(B, x)$ be the cluster algebra with trivial coefficients, where (B, x) is the initial seed [FZ2]. Let $\mathbf{1} = \{1\}$ be the *trivial semifield* and $\pi_{\mathbf{1}} : \mathbb{Q}_{\mathrm{sf}}(y) \to \mathbf{1}$, $y_{\mathbf{i}} \mapsto 1$ be the projection. Let $[x_{\mathbf{i}}(u)]_{\mathbf{1}}$ denote the image of $x_{\mathbf{i}}(u)$ by the algebra homomorphism $\mathcal{A}(B, x, y) \to \mathcal{A}(B, x)$ induced from $\pi_{\mathbf{1}}$. It is called the *trivial evaluation*.

Recall that $G(b, k, v; a, m, u)$ is defined in (2.10).

Lemma 3.4. *The family* $\{x^{(a)}_m(u) \mid (a,m,u) \in \mathcal{I}_{\ell+}\}$ *satisfies a system of relations*

$$\begin{aligned} x^{(a)}_m\left(u-\tfrac{1}{t_a}\right) x^{(a)}_m\left(u+\tfrac{1}{t_a}\right) = & \frac{y^{(a)}_m(u)}{1+y^{(a)}_m(u)} \prod_{(b,k,v)\in\mathcal{I}_{\ell+}} x^{(b)}_k(v)^{G(b,k,v;\,a,m,u)} \\ & + \frac{1}{1+y^{(a)}_m(u)} x^{(a)}_{m-1}(u) x^{(a)}_{m+1}(u), \end{aligned} \tag{3.16}$$

where $(a,m,u) \in \mathcal{I}'_{\ell+}$. *In particular, the family* $\{[x^{(a)}_m(u)]_{\mathbf{1}} \mid (a,m,u) \in \mathcal{I}_{\ell+}\}$ *satisfies the T-system* $\mathbb{T}_\ell(C_r)$ *in* $\mathcal{A}(B,x)$ *by replacing* $T^{(a)}_m(u)$ *with* $[x^{(a)}_m(u)]_{\mathbf{1}}$.

Definition 3.5. The *T-subalgebra* $\mathcal{A}_T(B,x)$ *of* $\mathcal{A}(B,x,y)$ *associated with the sequence* (3.11) is the subalgebra of $\mathcal{A}(B,x)$ generated by $[x_{\mathbf{i}}(u)]_{\mathbf{1}}$ ($(\mathbf{i},u) \in \mathbf{I} \times \frac{1}{2}\mathbb{Z}$).

Theorem 3.6. *The ring* $\mathcal{T}^\circ_\ell(C_r)_+$ *is isomorphic to* $\mathcal{A}_T(B,x)$ *by the correspondence* $T^{(a)}_m(u) \mapsto [x^{(a)}_m(u)]_{\mathbf{1}}$.

3.5. **Y-system and cluster algebra.** The result in this subsection is completely parallel to the B_r case [IIKKN].

Lemma 3.7. *The family* $\{y^{(a)}_m(u) \mid (a,m,u) \in \mathcal{I}'_{\ell+}\}$ *satisfies the Y-system* $\mathbb{Y}_\ell(C_r)$ *by replacing* $Y^{(a)}_m(u)$ *with* $y^{(a)}_m(u)$.

Definition 3.8. The *Y-subgroup* $\mathcal{G}_Y(B,y)$ *of* $\mathcal{G}(B,y)$ *associated with the sequence* (3.11) is the subgroup of $\mathcal{G}(B,y)$ generated by $y_{\mathbf{i}}(u)$ ($(\mathbf{i},u) \in \mathbf{I} \times \frac{1}{2}\mathbb{Z}$) and $1+y_{\mathbf{i}}(u)$ ($(\mathbf{i},u) : \mathbf{p}_+$ or $\mathbf{p}_-$).

Theorem 3.9. *The group* $\mathcal{Y}^\circ_\ell(C_r)_+$ *is isomorphic to* $\mathcal{G}_Y(B,y)$ *by the correspondence* $Y^{(a)}_m(u) \mapsto y^{(a)}_m(u)$ *and* $1+Y^{(a)}_m(u) \mapsto 1+y^{(a)}_m(u)$.

3.6. **Tropical Y-system at level 2.** The *tropical semifield* Trop(y) is an abelian multiplicative group freely generated by the elements $y_{\mathbf{i}}$ ($\mathbf{i} \in \mathbf{I}$) with the addition $\oplus$

$$\prod_{\mathbf{i}\in\mathbf{I}} y_{\mathbf{i}}^{a_{\mathbf{i}}} \oplus \prod_{\mathbf{i}\in\mathbf{I}} y_{\mathbf{i}}^{b_{\mathbf{i}}} = \prod_{\mathbf{i}\in\mathbf{I}} y_{\mathbf{i}}^{\min(a_{\mathbf{i}},b_{\mathbf{i}})}. \tag{3.17}$$

Let $\pi_{\mathbf{T}} : \mathbb{Q}_{\mathrm{sf}}(y) \to \mathrm{Trop}(y)$, $y_{\mathbf{i}} \mapsto y_{\mathbf{i}}$ be the projection. Let $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ and $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ denote the images of $y_{\mathbf{i}}(u)$ and $\mathcal{G}_Y(B,y)$ by the multiplicative group homomorphism induced from $\pi_{\mathbf{T}}$, respectively. They are called the *tropical evaluations*, and the resulting relations in the group $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ is called the *tropical Y-system.*

We say a (Laurent) monomial $m = \prod_{\mathbf{i}\in\mathbf{I}} y_{\mathbf{i}}^{k_{\mathbf{i}}}$ is *positive* (resp. *negative*) if $m \neq 1$ and $k_{\mathbf{i}} \geq 0$ (resp. $k_{\mathbf{i}} \leq 0$) for any $\mathbf{i}$.

The following properties of the tropical Y-system at level 2 will be the key in the entire method.

Proposition 3.10. *For* $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ *with* $B = B_2(C_r)$, *the following facts hold.*

(i) Let u *be in the region* $0 \leq u < 2$. *For any* $(\mathbf{i},u) : \mathbf{p}_+$, *the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is positive.*

(ii) Let u *be in the region* $-h^\vee \leq u < 0$.

- *(a) Let* $\mathbf{i} = (i,2)$ $(i \leq r-1)$, $(r,1)$, *or* $(r+1,1)$. *For any* $(\mathbf{i},u) : \mathbf{p}_+$, *the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is negative.*
- *(b) Let* $\mathbf{i} = (i,1), (i,3)$ $(i \leq r-1)$. *For any* $(\mathbf{i},u) : \mathbf{p}_+$, *the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is positive for* $u = -\frac{1}{2}h^\vee, -\frac{1}{2}h^\vee - \frac{1}{2}$ *and negative otherwise.*

(iii) $y_{ii'}(2) = y^{-1}_{i,4-i'}$ *if* $i \le r-1$ *and* $y^{-1}_{ii'}$ *if* $i = r, r+1$.

(iv) For even r, $y_{ii'}(-h^\vee) = y^{-1}_{ii'}$ *if* $i \le r-1$ *and* $y^{-1}_{2r+1-i,i'}$ *if* $i = r, r+1$. *For odd* r, $y_{ii'}(-h^\vee) = y^{-1}_{ii'}$.

One can directly verify (i) and (iii) in the same way as the B_r case [IIKKN, Proposition 3.2]. In the rest of this subsection we give the outline of the proof of (ii) and (iv). Note that (ii) and (iv) can be proved independently for each variable $y_{\mathbf{i}}$. (To be precise, we also need to assure that each monomial is not 1 in total. However, this can be easily followed up, so that we do not describe details here.) Below we separate the variables into two parts. Here is a brief summary of the results.

(1) *The D part.* The powers of $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ in the variables $y_{i,2}$ $(i \le r-1)$ and $y_{r,1}$, $y_{r+1,1}$ are described by the root system of type D_{r+1} with a Coxeter-like transformation. It turns out that they are further described by (a subset of) the root system of type A_{2r+1} with the Coxeter transformation.

(2) *The A part.* The powers of $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ in the variables $y_{i,1}$ and $y_{i,3}$ $(i \le r-1)$, are mainly described by the root system of type A_{r-1} with the Coxeter transformation.

3.6.1. *D part.* Let us consider the D part first. Let D_{r+1} be the Dynkin diagram of type D with index set $J = \{1, \dots, r+1\}$. We assign the sign $+/-$ to vertices of D_{r+1} (no sign for r and $r+1$) as inherited from $Q_2(C_r)$.





Let $\Pi = \{\alpha_1, \dots, \alpha_{r+1}\}$, $-\Pi$, Φ_+ be the set of the simple roots, the negative simple roots, the positive roots, respectively, of type D_{r+1}. Following [FZ1], we introduce the *piecewise-linear analogue* σ_i of the simple reflection s_i, acting on the set of the *almost positive roots* $\Phi_{\ge -1} = \Phi_+ \sqcup (-\Pi)$, by

$$\begin{aligned} \sigma_i(\alpha) &= s_i(\alpha), \quad \alpha \in \Phi_+, \\ \sigma_i(-\alpha_j) &= \begin{cases} \alpha_j & j = i, \\ -\alpha_j & \text{otherwise.} \end{cases} \end{aligned} \tag{3.18}$$

Let

$$\sigma_+ = \prod_{i \in J_+} \sigma_i, \quad \sigma_- = \prod_{i \in J_-} \sigma_i, \tag{3.19}$$

where $J_\pm$ is the set of the vertices of D_{r+1} with property $\pm$. We define σ as the composition

$$\sigma = \sigma_- \sigma_+ \sigma_{r+1} \sigma_- \sigma_+ \sigma_r. \tag{3.20}$$

Lemma 3.11. *The following facts hold.*

(I) Let r *be even.*

(i) For $i \le r-1$, $\sigma^k(-\alpha_i) \in \Phi_+$, $(1 \le k \le r/2)$, $\sigma^{r/2+1}(-\alpha_i) = -\alpha_i$.

(ii) For $i \le r-1$, $\sigma^k(\alpha_i) \in \Phi_+$, $(0 \le k \le r/2)$, $\sigma^{r/2+1}(\alpha_i) = \alpha_i$.

(iii) $\sigma^k(-\alpha_r) \in \Phi_+$, $(1 \le k \le r/2)$, $\sigma^{r/2+1}(-\alpha_r) = -\alpha_{r+1}$.

(iv) $\sigma^k(-\alpha_{r+1}) \in \Phi_+$, $(1 \le k \le r/2+1)$, $\sigma^{r/2+2}(-\alpha_{r+1}) = -\alpha_r$.

(v) The elements in Φ_+ *in (i)–(iv) exhaust the set* Φ_+*, thereby providing the orbit decomposition of* Φ_+ *by* σ.

(II) Let r *be odd.*

(i) For $i \in J_+$, $\sigma^k(-\alpha_i) \in \Phi_+$, $(1 \le k \le r+1)$, $\sigma^{r+2}(-\alpha_i) = -\alpha_i$, $\sigma^{(r+1)/2}(-\alpha_i) = \alpha_i$.

(ii) For $i \in J_-$, $\sigma^k(-\alpha_i) \in \Phi_+$, $(1 \le k \le r+1)$, $\sigma^{r+2}(-\alpha_i) = -\alpha_i$, $\sigma^{(r+3)/2}(-\alpha_i) = \alpha_i$.

(iii) $\sigma^k(-\alpha_r) \in \Phi_+$, $(1 \le k \le (r+1)/2)$, $\sigma^{(r+3)/2}(-\alpha_r) = -\alpha_r$.

(iv) $\sigma^k(-\alpha_{r+1}) \in \Phi_+$, $(1 \le k \le (r+1)/2)$, $\sigma^{(r+3)/2}(-\alpha_{r+1}) = -\alpha_{r+1}$.

(v) The elements in Φ_+ *in (i)–(iv) exhaust the set* Φ_+*, thereby providing the orbit decomposition of* Φ_+ *by* σ.

Proof. They are verified by explicitly calculating $\sigma^k(-\alpha_i)$ and $\sigma^k(\alpha_i)$. The examples for $r = 10$ (for even r) and 9 (for odd r) are given in Tables 1 and 2, respectively, where we use the notations

(3.21)
$$\begin{aligned}&[i,j] = \alpha_i + \cdots + \alpha_j \quad (1 \le i < j \le r), \quad [i] = \alpha_i \quad (1 \le i \le r),\\ &\{i,j\} = (\alpha_i + \cdots + \alpha_{r-1}) + (\alpha_j + \cdots + \alpha_{r+1}) \quad (1 \le i < j \le r+1, i \le r-1),\end{aligned}$$

and $\{r+1\} = \alpha_{r+1}$. In fact, it is not difficult to read off the general rule from these examples. □

The orbits $\sigma(-\alpha_i)$ and $\sigma(\alpha_i)$ are further described by (a subset of) the *root system of type* A_{2r+1}. Let $\Pi' = \{\alpha'_1, \dots, \alpha'_{2r+1}\}$ and Φ'_+ be the sets of the simple roots and the positive roots of type A_{2r+1}, respectively, with standard index set $J' = \{1, \dots, 2r+1\}$. Define, $J'_+ = \{i \in J' \mid i - r \text{ is even}\}$, $J'_- = \{i \in J' \mid i - r \text{ is odd}\}$. We introduce the notations $[i,j]' = \alpha'_i + \cdots + \alpha'_j$ $(1 \le i < j \le 2r+1)$ and $[i]' = \alpha'_i$ as parallel to (3.21). Let $O'_i = \{(\sigma')^k(-\alpha'_i) \mid 1 \le k \le r+1\}$ be the orbit of $-\alpha'_i$ in Φ'_+ by $\sigma' = \sigma'_-\sigma'_+$, $\sigma'_\pm = \prod_{i \in J'_\pm} \sigma'_i$, where σ'_i is the piecewise-linear analogue of the simple reflection s'_i as (3.18).

Lemma 3.12. *Let*

$$\rho : \Phi_+ \to \bigsqcup_{i=1}^{r} O'_i \tag{3.22}$$

be the map defined by

$$\begin{aligned}&[i,j] \mapsto \begin{cases}[i,j]' & j-r\text{: odd}\\ [2r+2-j, 2r+2-i]' & j-r\text{: even,}\end{cases}\\ &\{i,j\} \mapsto \begin{cases}[i, 2r+2-j]' & j-r\text{: odd}\\ [j, 2r+2-i]' & j-r\text{: even,}\end{cases}\\ &\{r+1\} \mapsto [r, r+1]',\end{aligned} \tag{3.23}$$

where $[i] = [i,i]$*. Then,* ρ *is a bijection. Furthermore, under the bijection* ρ*, the action of* σ *is translated into the one of the square of the Coxeter element* $s' = s'_-s'_+$ *of type* A_{2r+1} *acting on* Φ'_+*, where* $s'_\pm = \prod_{i \in J'_\pm} s'_i$.

		-1	-2	-3	-4	-5	-6	-7	-8	-9	-10	-11	
1 $-$	$-\alpha_1$	$[1]$	$[2,3]$	$[4,5]$	$[6,7]$	$[8,9]$	$\{11\}$	$[9,10]$	$[7,8]$	$[5,6]$	$[3,4]$	$[1,2]$	$-\alpha_1$ α_1
2 $+$	α_2 $-\alpha_2$	$[1,3]$	$[2,5]$	$[4,7]$	$[6,9]$	$\{8,11\}$	$\{9,10\}$	$[7,10]$	$[5,8]$	$[3,6]$	$[1,4]$	$[2]$	$-\alpha_2$
3 $-$	$-\alpha_3$	$[3]$	$[1,5]$	$[2,7]$	$[4,9]$	$\{6,11\}$	$\{8,9\}$	$\{7,10\}$	$[5,10]$	$[3,8]$	$[1,6]$	$[2,4]$	$-\alpha_3$ α_3
4 $+$	α_4 $-\alpha_4$	$[3,5]$	$[1,7]$	$[2,9]$	$\{4,11\}$	$\{6,9\}$	$\{7,8\}$	$\{5,10\}$	$[3,10]$	$[1,8]$	$[2,6]$	$[4]$	$-\alpha_4$
5 $-$	$-\alpha_5$	$[5]$	$[3,7]$	$[1,9]$	$\{2,11\}$	$\{4,9\}$	$\{6,7\}$	$\{5,8\}$	$\{3,10\}$	$[1,10]$	$[2,8]$	$[4,6]$	$-\alpha_5$ α_5
6 $+$	α_6 $-\alpha_6$	$[5,7]$	$[3,9]$	$\{1,11\}$	$\{2,9\}$	$\{4,7\}$	$\{5,6\}$	$\{3,8\}$	$\{1,10\}$	$[2,10]$	$[4,8]$	$[6]$	$-\alpha_6$
7 $-$	$-\alpha_7$	$[7]$	$[5,9]$	$\{3,11\}$	$\{1,9\}$	$\{2,7\}$	$\{4,5\}$	$\{3,6\}$	$\{1,8\}$	$\{2,10\}$	$[4,10]$	$[6,8]$	$-\alpha_7$ α_7
8 $+$	α_8 $-\alpha_8$	$[7,9]$	$\{5,11\}$	$\{3,9\}$	$\{1,7\}$	$\{2,5\}$	$\{3,4\}$	$\{1,6\}$	$\{2,8\}$	$\{4,10\}$	$[6,10]$	$[8]$	$-\alpha_8$
9 $-$	$-\alpha_9$	$[9]$	$\{7,11\}$	$\{5,9\}$	$\{3,7\}$	$\{1,5\}$	$\{2,3\}$	$\{1,4\}$	$\{2,6\}$	$\{4,8\}$	$\{6,10\}$	$[8,10]$	$-\alpha_9$ α_9
	$-\alpha_{11}$ $-\alpha_{10}$	$\{9,11\}$	$\{7,9\}$	$\{5,7\}$	$\{3,5\}$	$\{1,3\}$	$\{1,2\}$	$\{2,4\}$	$\{4,6\}$	$\{6,8\}$	$\{8,10\}$	$[10]$	$-\alpha_{11}$ $-\alpha_{10}$

TABLE 1. The orbits $\sigma^k(-\alpha_i)$ and $\sigma^k(\alpha_i)$ in Φ_+ by σ of (3.20) for $r = 10$. The orbits of $-\alpha_i$ and α_i $(i \leq 8)$, for example, $-\alpha_1 \to [2,3] \to [6,7] \to \cdots \to -\alpha_1$ and $\alpha_1 \to [4,5] \to [8,9] \to \cdots \to \alpha_1$, are aligned alternatively. The orbits of $-\alpha_{10}$ and $-\alpha_{11}$, namely, $-\alpha_{10} \to \{7,9\} \to \{3,5\} \to \cdots \to -\alpha_{11}$, and $-\alpha_{11} \to \{9,11\} \to \{5,7\} \to \cdots \to -\alpha_{10}$, are aligned alternatively. The numbers $-1, -2, \cdots$ in the head line will be identified with the parameter u in (3.24).

			-1		-2		-3		-4		-5		-6		-7		-8		-9		-10	
1 +	α_1 $-\alpha_1$		[1,2]		[3,4]		[5,6]		[7,8]		{10}		[8,9]		[6,7]		[4,5]		[2,3]		[1]	$-\alpha_1$
2 −	$-\alpha_2$	[2]		[1,4]		[3,6]		[5,8]		{7,10}		{8, 9}		[6,9]		[4,7]		[2,5]		[1, 3]		$-\alpha_2$ α_2
3 +	α_3 $-\alpha_3$		[2,4]		[1,6]		[3,8]		{5,10}		{7, 8}		{6, 9}		[4,9]		[2,7]		[1,5]		[3]	$-\alpha_3$
4 −	$-\alpha_4$	[4]		[2,6]		[1,8]		{3,10}		{5, 8}		{6, 7}		{4, 9}		[2,9]		[1,7]		[3, 5]		$-\alpha_4$ α_4
5 +	α_5 $-\alpha_5$		[4,6]		[2,8]		{1,10}		{3, 8}		{5, 6}		{4, 7}		{2, 9}		[1,9]		[3,7]		[5]	$-\alpha_5$
6 −	$-\alpha_6$	[6]		[4,8]		{2,10}		{1, 8}		{3, 6}		{4, 5}		{2, 7}		{1, 9}		[3,9]		[5, 7]		$-\alpha_6$ α_6
7 +	α_7 $-\alpha_7$		[6,8]		{4,10}		{2, 8}		{1, 6}		{3, 4}		{2, 5}		{1, 7}		{3, 9}		[5,9]		[7]	$-\alpha_7$
8 −	$-\alpha_8$	[8]		{6,10}		{4, 8}		{2, 6}		{1, 4}		{2, 3}		{1, 5}		{3, 7}		{5, 9}		[7,9]		$-\alpha_8$ α_8
	$-\alpha_{10}$ $-\alpha_9$		{8,10}		{6, 8}		{4, 6}		{2, 4}		{1, 2}		{1, 3}		{3, 5}		{5, 7}		{7, 9}		[9]	$-\alpha_{10}$ $-\alpha_9$

TABLE 2. The orbit of $\sigma^k(-\alpha_i)$ in Φ_+ by σ of (3.20) for $r = 9$. The orbit of $-\alpha_i$ $(i \leq 8)$, for example, $-\alpha_1 \to [3,4] \to [7,8] \to \cdots \to \alpha_1 \to [1,2] \to [5,6] \to \cdots \to -\alpha_1$, is aligned in a cyclic and alternative way. The orbits of $-\alpha_9$ and $-\alpha_{10}$, namely, $-\alpha_9 \to \{6,8\} \to \{2,4\} \to \cdots \to -\alpha_9$, and $-\alpha_{10} \to \{8,10\} \to \{4,6\} \to \cdots \to -\alpha_{10}$, are aligned alternatively. The numbers $-1, -2, \cdots$ in the head line will be identified with the parameter u in (3.24).

For $-h^\vee \le u < 0$, define

$$\alpha_i(u) = \begin{cases} \sigma^{-u/2}(-\alpha_i) & i \in J_+,\ u \equiv 0, \\ \sigma^{-(u-1)/2}(\alpha_i) & i \in J_+,\ u \equiv -1, \\ \sigma^{-(2u-1)/4}(-\alpha_i) & i \in J_-,\ u \equiv -\frac{3}{2}, \\ \sigma^{-(2u+1)/4}(\alpha_i) & i \in J_-,\ u \equiv -\frac{1}{2}, \\ \sigma^{-u/2}(-\alpha_r) & i = r,\ u \equiv 0, \\ \sigma^{-(u-1)/2}(-\alpha_{r+1}) & i = r+1,\ u \equiv -1, \end{cases} \tag{3.24}$$

where $\equiv$ is modulo $2\mathbb{Z}$. Note that they correspond to the positive roots in Tables 1 and 2 with u being the parameter in the head lines. By Lemma 3.11 they are all the positive roots of D_{r+1}.

Lemma 3.13. *The family in* (3.24) *satisfies the recurrence relations*

$$\begin{gathered} \alpha_i(u-\tfrac{1}{2}) + \alpha_i(u+\tfrac{1}{2}) = \alpha_{i-1}(u) + \alpha_{i+1}(u) \quad (1 \le i \le r-2), \\ \alpha_{r-1}(u-\tfrac{1}{2}) + \alpha_{r-1}(u+\tfrac{1}{2}) = \begin{cases} \alpha_{r-2}(u) + \alpha_r(u) & (u\text{: even}) \\ \alpha_{r-2}(u) + \alpha_{r+1}(u) & (u\text{: odd}), \end{cases} \\ \alpha_r(u-1) + \alpha_r(u+1) = \alpha_{r-1}(u-\tfrac{1}{2}) + \alpha_{r-1}(u+\tfrac{1}{2}) \quad (u\text{: odd}), \\ \alpha_{r+1}(u-1) + \alpha_{r+1}(u+1) = \alpha_{r-1}(u-\tfrac{1}{2}) + \alpha_{r-1}(u+\tfrac{1}{2}) \quad (u\text{: even}), \end{gathered} \tag{3.25}$$

where $\alpha_0(u) = 0$.

Proof. These relations are easily verified by the explicit expressions of $\alpha_i(u)$. See Tables 1 and 2. The first two relations are also obtained from Lemma 3.12 and [FZ2, Eq. (10.9)]. □

Let us return to prove (ii) of Proposition 3.10 for the D part. For a monomial m in $y = (y_{\mathbf{i}})_{\mathbf{i}\in\mathbf{I}}$, let $\pi_D(m)$ denote the specialization with $y_{i,1} = y_{i,3} = 1$ $(i \le r-1)$. For simplicity, we set $y_{i2} = y_i$ $(i \le r-1)$, $y_{r1} = y_r$, $y_{r+1,1} = y_{r+1}$, and also, $y_{i2}(u) = y_i(u)$ $(i \le r-1)$, $y_{r1}(u) = y_r(u)$, $y_{r+1,1}(u) = y_r(u)$. We define the vectors $\mathbf{t}_i(u) = (t_i(u)_k)_{k=1}^{r+1}$ by

$$\pi_D([y_i(u)]_{\mathbf{T}}) = \prod_{k=1}^{r+1} y_k^{t_i(u)_k}. \tag{3.26}$$

We also identify each vector $\mathbf{t}_i(u)$ with $\alpha = \sum_{k=1}^{r+1} t_i(u)_k \alpha_k \in \mathbb{Z}\Pi$.

Proposition 3.14. *Let* $-h^\vee \le u < 0$. *Then, we have*

$$\mathbf{t}_i(u) = -\alpha_i(u) \tag{3.27}$$

for (i,u) *in* (3.24), *and*

$$\pi_D([y_{i1}(u)]_{\mathbf{T}}) = \pi_D([y_{i3}(u)]_{\mathbf{T}}) = 1 \quad (i \le r-1,\ r+i+2u\text{: even}). \tag{3.28}$$

Note that these formulas determine $\pi_D([y_{\mathbf{i}}(u)]_{\mathbf{T}})$ for any $(\mathbf{i}, u) : \mathbf{p}_+$.

Proof. We can verify the claim for $-2 \le u \le -\frac{1}{2}$ by direct computation. Then, by induction on u in the backward direction, one can establish the claim, together with the recurrence relations among $\mathbf{t}_i(u)$'s with (i,u) in (3.24),

$$\begin{aligned}
&\mathbf{t}_i(u-\tfrac{1}{2})+\mathbf{t}_i(u+\tfrac{1}{2})=\mathbf{t}_{i-1}(u)+\mathbf{t}_{i+1}(u)\quad(1\le i\le r-2),\\
&\mathbf{t}_{r-1}(u-\tfrac{1}{2})+\mathbf{t}_{r-1}(u+\tfrac{1}{2})=\begin{cases}\mathbf{t}_{r-2}(u)+\mathbf{t}_r(u) & (u\text{: even})\\ \mathbf{t}_{r-2}(u)+\mathbf{t}_{r+1}(u) & (u\text{: odd}),\end{cases}\\
&\mathbf{t}_r(u-1)+\mathbf{t}_r(u+1)=\mathbf{t}_{r-1}(u-\tfrac{1}{2})+\mathbf{t}_{r-1}(u+\tfrac{1}{2})\quad(u\text{: odd}),\\
&\mathbf{t}_{r+1}(u-1)+\mathbf{t}_{r+1}(u+1)=\mathbf{t}_{r-1}(u-\tfrac{1}{2})+\mathbf{t}_{r-1}(u+\tfrac{1}{2})\quad(u\text{: even}).
\end{aligned}\tag{3.29}$$

Note that (3.29) coincides with (3.25) under (3.27). To derive (3.29), one uses the mutations as in [IIKKN, Figure 6] (or the tropical version of the Y-system $\mathbb{Y}_2(C_r)$ directly) and the positivity/negativity of $\pi_D([y_{\mathbf{i}}(u)]_{\mathbf{T}})$ resulting from (3.27) and (3.28) by induction hypothesis. □

Now (ii) and (iv) in Proposition 3.10 for the D part follow from Proposition 3.14.

3.6.2. *A part.* The A part can be studied in a similar way to the D part. Thus, we present only the result.

First we note that the quiver $Q_2(C_r)$ is symmetric under the exchange $y_{i,1}\leftrightarrow y_{i,3}$ $(i\le r-1)$. Thus, one can concentrate on the powers of $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ in the variables $y_{i,1}$ $(i\le r-1)$.

Let A_{r-1} be the Dynkin diagram of type A with index set $J=\{1,\dots,r-1\}$. We assign the sign $+/-$ to vertices (except for r) of A_{r-1} as inherited from $Q_2(C_r)$.



r: even



r: odd

Let $\Pi=\{\alpha_1,\dots,\alpha_{r-1}\}$, $-\Pi$, Φ_+ be the set of the simple roots, the negative simple roots, the positive roots, respectively, of type A_{r-1}. Again, we introduce the piecewise-linear analogue σ_i of the simple reflection s_i, acting on $\Phi_{\ge -1}=\Phi_+\sqcup(-\Pi)$ as (3.18). Let

$$\sigma_+=\prod_{i\in J_+}\sigma_i,\quad \sigma_-=\prod_{i\in J_-}\sigma_i,\tag{3.30}$$

where $J_\pm$ is the set of the vertices of A_{r-1} with property $\pm$. We define σ as the composition

$$\sigma=\sigma_-\sigma_+.\tag{3.31}$$

For a monomial m in $y=(y_{\mathbf{i}})_{\mathbf{i}\in\mathbf{I}}$, let $\pi_A(m)$ denote the specialization with $y_{i,2}=y_{i,3}=1$ $(i\le r-1)$ and $y_{r1}=y_{r+1,1}=1$. We set $y_{i1}=y_i$ $(i\le r-1)$.

We define the vectors $\mathbf{t}_{\mathbf{i}}(u)=(t_{\mathbf{i}}(u)_k)_{k=1}^{r-1}$ by

$$\pi_A([y_{\mathbf{i}}(u)]_{\mathbf{T}})=\prod_{k=1}^{r-1}y_k^{t_{\mathbf{i}}(u)_k}.\tag{3.32}$$

We also identify each vector $\mathbf{t}_{\mathbf{i}}(u)$ with $\alpha=\sum_{k=1}^{r-1}t_{\mathbf{i}}(u)_k\alpha_k\in\mathbb{Z}\Pi$.

With these notations, the result for the A part is summarized as follows.

Proposition 3.15. *Let $-h^\vee \le u < 0$. For $(\mathbf{i},u) : \mathbf{p}_+$, $\mathbf{t}_{\mathbf{i}}(u)$ is given by, for $i \le r-1$,*

$$\begin{aligned}
&\mathbf{t}_{i1}(u) = \begin{cases} -\sigma^{-u}(-\alpha_i) & i \in J_+ \\ -\sigma^{-(2u-1)/2}(-\alpha_i) & i \in J_-, \end{cases}\\
&\mathbf{t}_{i2}(u) = \begin{cases} -[2r+2-i+2u, r-1] & -\tfrac{1}{2}h^\vee \le u < 0 \\ -[-1-i-2u, r-1] & -h^\vee \le u < -\tfrac{1}{2}h^\vee, \end{cases}\\
&\mathbf{t}_{i3}(u) = 0,
\end{aligned} \tag{3.33}$$

and

$$\begin{aligned}
&\mathbf{t}_{r1}(u) = \begin{cases} -[r+2+2u, r-1] & -\tfrac{1}{2}h^\vee \le u < 0 \\ -[-1-r-2u, r-1] & -h^\vee \le u < -\tfrac{1}{2}h^\vee, \end{cases}\\
&\mathbf{t}_{r+1,1}(u) = \begin{cases} -[r+2+2u, r-1] & -\tfrac{1}{2}h^\vee \le u < 0 \\ -[-1-r-2u, r-1] & -h^\vee \le u < -\tfrac{1}{2}h^\vee, \end{cases}
\end{aligned} \tag{3.34}$$

where $[i,j] = \alpha_i + \cdots + \alpha_j$ if $i \le j$ and 0 if $i > j$.

Note that $\mathbf{t}_{i1}(-\frac{h^\vee}{2}) = \alpha_{r-i}$ ($i \in J_-$ for even r and $i \in J_+$ for odd r) and $\mathbf{t}_{i1}(-\frac{h^\vee}{2} - \frac{1}{2}) = \alpha_{r-i}$ ($i \in J_+$ for even r and $i \in J_-$ for odd r), and that they are the only positive monomials in (3.33) and (3.34). Now (ii) and (iv) in Proposition 3.10 for the A part follow from Proposition 3.15.

This completes the proof of Proposition 3.10.

3.7. Tropical Y-systems at higher levels.

By the same method for the B_r case [IIKKN, Proposition 4.1], one can establish the 'factorization property' of the tropical Y-system at higher levels. As a result, we obtain a generalization of Proposition 3.10.

Proposition 3.16. *For $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ with $B = B_\ell(C_r)$, the following facts hold.*

(i) Let u be in the region $0 \le u < \ell$. For any $(\mathbf{i},u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is positive.

(ii) Let u be in the region $-h^\vee \le u < 0$.

- *(a) Let $\mathbf{i} \in \mathbf{I}^\circ$ or $\mathbf{i} = (i,i')$ ($i \le r-1, i' \in 2\mathbb{N}$). For any $(\mathbf{i},u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is negative.*
- *(b) Let $\mathbf{i} = (i,i')$ ($i \le r-1, i' \notin 2\mathbb{N}$). For any $(\mathbf{i},u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is positive for $u = -\frac{1}{2}h^\vee, -\frac{1}{2}h^\vee - \frac{1}{2}$ and negative otherwise.*

(iii) $y_{ii'}(\ell) = y^{-1}_{i,2\ell-i'}$ if $i \le r-1$ and $y^{-1}_{i,\ell-i'}$ if $i = r, r+1$.

(iv) For even r, $y_{ii'}(-h^\vee) = y^{-1}_{ii'}$ if $i \le r-1$ and $y^{-1}_{2r+1-i,i'}$ if $i = r, r+1$. For odd r, $y_{ii'}(-h^\vee) = y^{-1}_{ii'}$.

We obtain two important corollaries of Propositions 3.10 and 3.16.

Theorem 3.17. *For $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ the following relations hold.*

(i) Half periodicity: $[y_{\mathbf{i}}(u + h^\vee + \ell)]_{\mathbf{T}} = [y_{\boldsymbol{\omega}(\mathbf{i})}(u)]_{\mathbf{T}}$.

(ii) Full periodicity: $[y_{\mathbf{i}}(u + 2(h^\vee + \ell))]_{\mathbf{T}} = [y_{\mathbf{i}}(u)]_{\mathbf{T}}$.

Theorem 3.18. *For $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$, let N_+ and N_- denote the total numbers of the positive and negative monomials, respectively, among $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ for $(\mathbf{i},u) : \mathbf{p}_+$ in the*



FIGURE 4. The quiver $Q_\ell(F_4)$ for even ℓ (upper) and for odd ℓ (lower), where we identify the right column in the left quiver with the middle column in the right quiver.

region $0 \leq u < 2(h^\vee + \ell)$. *Then, we have*

$$N_+ = 2\ell(2r\ell - \ell - 1), \quad N_- = 2r(2\ell r - r - 1). \tag{3.35}$$

We observe the symmetry (the *level-rank duality*) for the numbers N_+ and N_- under the exchange of r and ℓ.

3.8. **Periodicities and dilogarithm identities.** Applying [IIKKN, Theorem 5.1] to Theorem 3.17, we obtain the periodicities:

Theorem 3.19. *For $\mathcal{A}(B,x,y)$, the following relations hold.*
(i) Half periodicity: $x_{\mathbf{i}}(u + h^\vee + \ell) = x_{\boldsymbol{\omega}(\mathbf{i})}(u)$.
(ii) Full periodicity: $x_{\mathbf{i}}(u + 2(h^\vee + \ell)) = x_{\mathbf{i}}(u)$.

Theorem 3.20. *For $\mathcal{G}(B,y)$, the following relations hold.*
(i) Half periodicity: $y_{\mathbf{i}}(u + h^\vee + \ell) = y_{\boldsymbol{\omega}(\mathbf{i})}(u)$.
(ii) Full periodicity: $y_{\mathbf{i}}(u + 2(h^\vee + \ell)) = y_{\mathbf{i}}(u)$.

Then, Theorems 2.5 and 2.6 for C_r follow from Theorems 3.6, 3.9, 3.19, and 3.20. Furthermore, Theorem 2.10 for C_r is obtained from the above periodicities and Theorem 3.18 as in the B_r case [IIKKN, Section 6].

4. TYPE F_4

The F_4 case is quite parallel to the B_r and C_r case. We do not repeat the same definitions unless otherwise mentioned. Again, the properties of the tropical Y-system at level 2 (Proposition 4.7) are crucial and specific to F_4.

4.1. **Parity decompositions of T and Y-systems.** For a triplet $(a,m,u)\in\mathcal{I}_\ell$, we reset the 'parity conditions' $\mathbf{P}_+$ and $\mathbf{P}_-$ by

$$\mathbf{P}_+:\ 2u \text{ is even if } a=1,2;\ a+m+2u \text{ is odd if } a=3,4, \tag{4.1}$$

$$\mathbf{P}_-:\ 2u \text{ is odd if } a=1,2;\ a+m+2u \text{ is even if } a=3,4. \tag{4.2}$$

Then, we have $\mathcal{T}^\circ_\ell(F_4)_+\simeq\mathcal{T}^\circ_\ell(F_4)_-$ by $T^{(a)}_m(u)\mapsto T^{(a)}_m(u+\frac{1}{2})$ and

$$\mathcal{T}^\circ_\ell(F_4)\simeq\mathcal{T}^\circ_\ell(F_4)_+\otimes_{\mathbb{Z}}\mathcal{T}^\circ_\ell(F_4)_-. \tag{4.3}$$

For a triplet $(a,m,u)\in\mathcal{I}_\ell$, we reset the parity condition $\mathbf{P}'_+$ and $\mathbf{P}'_-$ by

$$\mathbf{P}'_+:\ 2u \text{ is even if } a=1,2;\ a+m+2u \text{ is even if } a=3,4, \tag{4.4}$$

$$\mathbf{P}'_-:\ 2u \text{ is odd if } a=1,2;\ a+m+2u \text{ is odd if } a=3,4. \tag{4.5}$$

We have

$$(a,m,u):\mathbf{P}'_+\iff(a,m,u\pm\tfrac{1}{t_a}):\mathbf{P}_+. \tag{4.6}$$

Also, we have $\mathcal{Y}^\circ_\ell(F_4)_+\simeq\mathcal{Y}^\circ_\ell(F_4)_-$ by $Y^{(a)}_m(u)\mapsto Y^{(a)}_m(u+\frac{1}{2})$, $1+Y^{(a)}_m(u)\mapsto 1+Y^{(a)}_m(u+\frac{1}{2})$, and

$$\mathcal{Y}^\circ_\ell(F_4)\simeq\mathcal{Y}^\circ_\ell(F_4)_+\times\mathcal{Y}^\circ_\ell(F_4)_-. \tag{4.7}$$

4.2. **Quiver $Q_\ell(F_4)$.** With type F_4 and $\ell\geq 2$ we associate the quiver $Q_\ell(F_4)$ by Figure 4, where the right column in the left quiver and the middle column in the right quiver are identified. Also, we assign the empty or filled circle ○/● and the sign $+/-$ to each vertex.

Let us choose the index set $\mathbf{I}$ of the vertices of $Q_\ell(F_4)$ so that $\mathbf{i}=(i,i')\in\mathbf{I}$ represents the vertex at the i'th row (from the bottom) of the ith column (from the left) in the right quiver for $i=1,2,3$, the one of the $i-1$th column in the right quiver for $i=5,6$, and the one of the left column in the left quiver for $i=4$. Thus, $i=1,\dots,6$, and $i'=1,\dots,\ell-1$ if $i=1,2,5,6$ and $i'=1,\dots,2\ell-1$ if $i=3,4$.

Let $\boldsymbol{r}$ be the involution acting on $\mathbf{I}$ by the left-right reflection of the right quiver. Let $\boldsymbol{\omega}$ be the involution acting on $\mathbf{I}$ by the up-down reflection of the left quiver and the 180° rotation of the right quiver.

Lemma 4.1. *Let $Q=Q_\ell(F_4)$.*

(i) We have the same periodic sequence of mutations of quivers as (3.9).

(ii) $\boldsymbol{\omega}(Q)=Q$ if $h^\vee+\ell$ is even, and $\boldsymbol{\omega}(Q)=\boldsymbol{r}(Q)$ if $h^\vee+\ell$ is odd.

4.3. **Cluster algebra and alternative labels.** Let $B_\ell(F_4)$ be the corresponding skew-symmetric matrix to the quiver $Q_\ell(F_4)$. In the rest of the section, we set the matrix $B=(B_{\mathbf{ij}})_{\mathbf{i},\mathbf{j}\in\mathbf{I}}=B_\ell(F_4)$ unless otherwise mentioned.

Let $\mathcal{A}(B,x,y)$ be the cluster algebra with coefficients in the universal semifield $\mathbb{Q}_{\mathrm{sf}}(y)$, and the coefficient group $\mathcal{G}(B,y)$ associated with $\mathcal{A}(B,x,y)$.

In view of Lemma 4.1 we set $x(0)=x$, $y(0)=y$ and define clusters $x(u)=(x_{\mathbf{i}}(u))_{\mathbf{i}\in\mathbf{I}}$ $(u\in\frac{1}{2}\mathbb{Z})$ and coefficient tuples $y(u)=(y_{\mathbf{i}}(u))_{\mathbf{i}\in\mathbf{I}}$ $(u\in\frac{1}{2}\mathbb{Z})$ by the sequence of mutations (3.11).

For a pair $(\mathbf{i},u)\in\mathbf{I}\times\frac{1}{2}\mathbb{Z}$, we set the same parity condition $\mathbf{p}_+$ and $\mathbf{p}_-$ as (3.12). We have (3.13), and each $(\mathbf{i},u):\mathbf{p}_+$ is a mutation point of (3.11) in the forward direction of u, and each $(\mathbf{i},u):\mathbf{p}_-$ is so in the backward direction of u as before.


$x(0)$
$x(\frac{1}{2})$
$\mu^\bullet_+\mu^\circ_+\updownarrow$
$\mu^\bullet_-\updownarrow$
$x^{(3)}_7(-\frac{1}{2})$
$x^{(4)}_6(-\frac{1}{2})$
$x^{(3)}_5(-\frac{1}{2})$
$x^{(4)}_4(-\frac{1}{2})$
$x^{(3)}_3(-\frac{1}{2})$
$x^{(4)}_2(-\frac{1}{2})$
$x^{(3)}_1(-\frac{1}{2})$
$x^{(1)}_3(-1)$
$x^{(2)}_3(-1)$
$x^{(2)}_2(-1)$
$x^{(1)}_2(-1)$
$x^{(1)}_1(-1)$
$x^{(2)}_1(-1)$
$x^{(4)}_7(0)$
$x^{(3)}_6(0)$
$x^{(4)}_5(0)$
$x^{(3)}_4(0)$
$x^{(4)}_3(0)$
$x^{(3)}_2(0)$
$x^{(4)}_1(0)$
$*$
$\diamond$

FIGURE 5. (Continues to Figure 6) Label of cluster variables $x_{\mathbf{i}}(u)$ by $\mathcal{I}_{\ell+}$ for F_4, $\ell = 4$. The middle column in the right quiver (marked by $\diamond$) is identified with the right column in the left quiver.

Lemma 4.2. *Below $\equiv$ means the equivalence modulo $2\mathbb{Z}$.*

(i) The map

$$\begin{aligned} g:\quad &\mathcal{I}_{\ell+} \quad\to\quad \{(\mathbf{i},u):\mathbf{p}_+\}\\ &(a,m,u-\tfrac{1}{t_a}) \mapsto \begin{cases} ((a,m),u) & a=1,2;\ a+m+u\equiv 0\\ ((7-a,m),u) & a=1,2;\ a+m+u\equiv 1\\ ((a,m),u) & a=3,4 \end{cases} \end{aligned} \tag{4.8}$$

is a bijection.

(ii) The map

$$\begin{aligned} g':\quad &\mathcal{I}'_{\ell+} \quad\to\quad \{(\mathbf{i},u):\mathbf{p}_+\}\\ &(a,m,u) \mapsto \begin{cases} ((a,m),u) & a=1,2;\ a+m+u\equiv 0\\ ((7-a,m),u) & a=1,2;\ a+m+u\equiv 1\\ ((a,m),u) & a=3,4 \end{cases} \end{aligned} \tag{4.9}$$



FIGURE 6. (Continues from Figure 5).

is a bijection.

We introduce alternative labels $x_{\mathbf{i}}(u) = x_m^{(a)}(u - 1/t_a)$ $((a, m, u - 1/t_a) \in \mathcal{I}_{\ell+})$ for $(\mathbf{i}, u) = g((a, m, u - 1/t_a))$ and $y_{\mathbf{i}}(u) = y_m^{(a)}(u)$ $((a, m, u) \in \mathcal{I}'_{\ell+})$ for $(\mathbf{i}, u) = g'((a, m, u))$, respectively. See Figures 5–6.

4.4. **T-system and cluster algebra.** The result in this subsection is completely parallel to the B_r and C_r cases.

Lemma 4.3. *The family* $\{x_m^{(a)}(u) \mid (a, m, u) \in \mathcal{I}_{\ell+}\}$ *satisfies the system of relations* (3.16) *with* $G(b, k, v; a, m, u)$ *for* $\mathbb{T}_\ell(F_4)$*. In particular, the family* $\{[x_m^{(a)}(u)]_{\mathbf{1}} \mid (a, m, u) \in \mathcal{I}_{\ell+}\}$ *satisfies the T-system* $\mathbb{T}_\ell(F_4)$ *in* $\mathcal{A}(B, x)$ *by replacing* $T_m^{(a)}(u)$ *with* $[x_m^{(a)}(u)]_{\mathbf{1}}$.

The T-subalgebra $\mathcal{A}_T(B, x)$ is defined as Definition 3.5.

Theorem 4.4. *The ring* $\mathcal{T}_\ell^\circ(F_4)_+$ *is isomorphic to* $\mathcal{A}_T(B, x)$ *by the correspondence* $T_m^{(a)}(u) \mapsto [x_m^{(a)}(u)]_{\mathbf{1}}$.

4.5. **Y-system and cluster algebra.** The result in this subsection is completely parallel to the B_r and C_r cases.

Lemma 4.5. *The family* $\{y^{(a)}_m(u) \mid (a,m,u) \in \mathcal{I}'_{\ell+}\}$ *satisfies the Y-system* $\mathbb{Y}_\ell(F_4)$ *by replacing* $Y^{(a)}_m(u)$ *with* $y^{(a)}_m(u)$.

The Y-subgroup $\mathcal{G}_Y(B,y)$ is defined as Definition 3.8.

Theorem 4.6. *The group* $\mathcal{Y}^\circ_\ell(F_4)_+$ *is isomorphic to* $\mathcal{G}_Y(B,y)$ *by the correspondence* $Y^{(a)}_m(u) \mapsto y^{(a)}_m(u)$ *and* $1+Y^{(a)}_m(u) \mapsto 1+y^{(a)}_m(u)$.

4.6. **Tropical Y-system at level 2.** By direct computations, the following properties are verified.

Proposition 4.7. *For* $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ *with* $B = B_2(F_4)$, *the following facts hold.*

(i) Let u *be in the region* $0 \le u < 2$. *For any* $(\mathbf{i},u) : \mathbf{p}_+$, *the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is positive.*

(ii) Let u *be in the region* $-h^\vee \le u < 0$.

- *(a) Let* $\mathbf{i} = (1,1),(2,1),(5,1),(6,1),(3,2)$, *or* $(4,2)$. *For any* $(\mathbf{i},u) : \mathbf{p}_+$, *the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is negative.*
- *(b) Let* $\mathbf{i} = (3,1),(3,3),(4,1)$, *or* $(4,3)$. *For any* $(\mathbf{i},u) : \mathbf{p}_+$, *the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is negative for* $u = -\frac{1}{2}, -1, -\frac{3}{2}, -3, -\frac{7}{2}, -4, -\frac{11}{2}, -6, -\frac{13}{2}, -8, -\frac{17}{2}, -9$ *and positive for* $u = -2, -\frac{5}{2}, -\frac{9}{2}, -5, -7, -\frac{15}{2}$.

(iii) $y_{ii'}(2) = y^{-1}_{ii'}$ *if* $i = 1,2,5,6$ *and* $y^{-1}_{i,4-i'}$ *if* $i = 3,4$.

(iv) $y_{ii'}(-h^\vee) = y^{-1}_{7-i,i'}$ *if* $i = 1,2,5,6$ *and* $y^{-1}_{ii'}$ *if* $i = 3,4$.

Also we have a description of the 'core part' of $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ in the region $-h^\vee \le u < 0$, corresponding to the D part for C_r, in terms of the root system of E_6. We use the following index of the Dynkin diagram E_6.



Let $\Pi = \{\alpha_1, \dots, \alpha_6\}$, $-\Pi$, Φ_+ be the set of the simple roots, the negative simple roots, the positive roots, respectively, of type E_6. Let σ_i be the piecewise-linear analogue of the simple reflection s_i, acting on the set of the almost positive roots $\Phi_{\ge -1} = \Phi_+ \sqcup (-\Pi)$. We write $\sum_i m_i\alpha_i \in \Phi_+$ as $[1^{m_1}, 2^{m_2}, \dots, 6^{m_6}]$; furthermore, $[1^0, 2^1, 3^1, 4^1, 5^0, 6^0]$, for example, is abbreviated as $[2,3,4]$.

We define σ as the composition

$$\sigma = \sigma_3(\sigma_4\sigma_2\sigma_6)\sigma_3(\sigma_4\sigma_1\sigma_5). \tag{4.10}$$

Lemma 4.8. *We have the orbits by σ*

(4.11)

$$\begin{aligned}
&-\alpha_1 \to [1,2,3] \to [2,3,4,5,6] \to [1,2,3^2,4,5] \to [5,6] \to -\alpha_6,\\
&-\alpha_2 \to [2,3] \to [1,2^2,3^2,4,5,6] \to [1,2^2,3^3,4^2,5^2,6]\\
&\qquad\qquad\qquad\qquad \to [1,2,3^2,4,5^2,6] \to [5] \to -\alpha_5,\\
&-\alpha_3 \to [2,3,4] \to [1,2,3^2,4,5,6] \to [2,3^2,4,5^2,6] \to [1,2,3,5] \to -\alpha_3,\\
&\ \ \alpha_3 \to [2,3,5,6] \to [1,2^2,3^2,4,5] \to [1,2,3,4,5,6] \to [3,4,5] \to \alpha_3,\\
&-\alpha_4 \to [2] \to [1,2,3,4] \to [3,4,5,6] \to [3,5] \to -\alpha_4,\\
&\ \ \alpha_4 \to [3,4] \to [3,5,6] \to [2,3,5] \to [1,2] \to \alpha_4,\\
&-\alpha_5 \to [2,3^2,4,5,6] \to [1,2^2,3^3,4,5^2,6] \to [1,2^2,3^2,4,5^2,6]\\
&\qquad\qquad\qquad\qquad \to [1,2,3,4,5] \to -\alpha_2,\\
&-\alpha_6 \to [6] \to [2,3^2,4,5] \to [1,2,3,5,6] \to [2,3,4,5] \to [1] \to -\alpha_1.
\end{aligned}$$

In particular, these elements in Φ_+ exhaust the set Φ_+, thereby providing the orbit decomposition of Φ_+ by σ.

For $-h^\vee \le u < 0$, define

$$\alpha_i(u) = \begin{cases} \sigma^{-u/2}(-\alpha_i) & i=1,4,5;\ u\equiv 0,\\ \sigma^{-(u-1)/2}(-\alpha_i) & i=2,6;\ u\equiv -1,\\ \sigma^{-(u-1)/2}(\alpha_4) & i=4;\ u\equiv -1,\\ \sigma^{-(2u-1)/4}(-\alpha_3) & i=3;\ u\equiv -\frac{3}{2},\\ \sigma^{-(2u+1)/4}(\alpha_3) & i=3;\ u\equiv -\frac{1}{2},\end{cases} \tag{4.12}$$

where $\equiv$ is mod $2\mathbb{Z}$. By Lemma 4.8 they are (all the) positive roots of E_6.

For a monomial m in $y = (y_{\mathbf{i}})_{\mathbf{i}\in\mathbf{I}}$, let $\pi_A(m)$ denote the specialization with $y_{31} = y_{33} = y_{41} = y_{43} = 1$. For simplicity, we set $y_{i1} = y_i$ $(i = 1,2,5,6)$, $y_{i2} = y_i$ $(i = 3,4)$, and also, $y_{i1}(u) = y_i(u)$ $(i = 1,2,5,6)$, $y_{i2}(u) = y_i(u)$ $(i = 3,4)$. We define the vectors $\mathbf{t}_i(u) = (t_i(u)_k)_{k=1}^6$ by

$$\pi_A([y_i(u)]_{\mathbf{T}}) = \prod_{k=1}^{6} y_k^{t_i(u)_k}. \tag{4.13}$$

We also identify each vector $\mathbf{t}_i(u)$ with $\alpha = \sum_{k=1}^6 t_i(u)_k\alpha_k \in \mathbb{Z}\Pi$.

Proposition 4.9. *Let $-h^\vee \le u < 0$. Then, we have*

$$\mathbf{t}_i(u) = -\alpha_i(u) \tag{4.14}$$

for (i,u) in (4.12).

4.7. **Tropical Y-systems at higher levels.**

Due to the factorization property, we obtain the following.

Proposition 4.10. *Let $\ell > 2$ be an integer. For $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ with $B = B_\ell(F_4)$, the following facts hold.*

(i) Let u be in the region $0 \le u < \ell$. For any $(\mathbf{i},u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is positive.

(ii) Let u be in the region $-h^\vee \le u < 0$.

(a) Let $\mathbf{i} \in \mathbf{I}^\circ$ *or* $\mathbf{i} = (3, i'), (4, i')$ $(i' \in 2\mathbb{N})$*. For any* $(\mathbf{i}, u) : \mathbf{p}_+$*, the monomial* $[y_\mathbf{i}(u)]_\mathbf{T}$ *is negative.*

(b) Let $\mathbf{i} = (3, i'), (4, i')$ $(i' \notin 2\mathbb{N})$*. For any* $(\mathbf{i}, u) : \mathbf{p}_+$*, the monomial* $[y_\mathbf{i}(u)]_\mathbf{T}$ *is negative for* $u = -\frac{1}{2}, -1, -\frac{3}{2}, -3, -\frac{7}{2}, -4, -\frac{11}{2}, -6, -\frac{13}{2}, -8, -\frac{17}{2}, -9$ *and positive for* $u = -2, -\frac{5}{2}, -\frac{9}{2}, -5, -7, -\frac{15}{2}$*.*

(iii) $y_{ii'}(\ell) = y^{-1}_{i,\ell-i'}$ *if* $i = 1, 2, 5, 6$ *and* $y^{-1}_{i,2\ell-i'}$ *if* $i = 3, 4$*.*

(iv) $y_{ii'}(-h^\vee) = y^{-1}_{7-i,i'}$ *if* $i = 1, 2, 5, 6$ *and* $y^{-1}_{ii'}$ *if* $i = 3, 4$*.*

We obtain corollaries of Propositions 4.7 and 4.10.

Theorem 4.11. *For* $[\mathcal{G}_Y(B, y)]_\mathbf{T}$ *the following relations hold.*

(i) Half periodicity: $[y_\mathbf{i}(u + h^\vee + \ell)]_\mathbf{T} = [y_{\boldsymbol{\omega}(\mathbf{i})}(u)]_\mathbf{T}$*.*

(ii) Full periodicity: $[y_\mathbf{i}(u + 2(h^\vee + \ell))]_\mathbf{T} = [y_\mathbf{i}(u)]_\mathbf{T}$*.*

Theorem 4.12. *For* $[\mathcal{G}_Y(B, y)]_\mathbf{T}$*, let* N_+ *and* N_- *denote the total numbers of the positive and negative monomials, respectively, among* $[y_\mathbf{i}(u)]_\mathbf{T}$ *for* $(\mathbf{i}, u) : \mathbf{p}_+$ *in the region* $0 \le u < 2(h^\vee + \ell)$*. Then, we have*

$$N_+ = 4\ell(3\ell + 1), \quad N_- = 24(4\ell - 3). \tag{4.15}$$

4.8. **Periodicities and dilogarithm identities.** Applying [IIKKN, Theorem 5.1] to Theorem 4.11, we obtain the periodicities:

Theorem 4.13. *For* $\mathcal{A}(B, x, y)$*, the following relations hold.*

(i) Half periodicity: $x_\mathbf{i}(u + h^\vee + \ell) = x_{\boldsymbol{\omega}(\mathbf{i})}(u)$*.*

(ii) Full periodicity: $x_\mathbf{i}(u + 2(h^\vee + \ell)) = x_\mathbf{i}(u)$*.*

Theorem 4.14. *For* $\mathcal{G}(B, y)$*, the following relations hold.*

(i) Half periodicity: $y_\mathbf{i}(u + h^\vee + \ell) = y_{\boldsymbol{\omega}(\mathbf{i})}(u)$*.*

(ii) Full periodicity: $y_\mathbf{i}(u + 2(h^\vee + \ell)) = y_\mathbf{i}(u)$*.*

Then, Theorems 2.5 and 2.6 for F_4 follow from Theorems 4.4, 4.6, 4.13, and 4.14. Furthermore, Theorem 2.10 for F_4 is obtained from the above periodicities and Theorem 4.12 as in the B_r case [IIKKN, Section 6].

5. TYPE G_2

The G_2 case is mostly parallel to the former cases, but slightly different because the number t in (2.2) is three. Again, the properties of the tropical Y-system at level 2 (Proposition 5.9) are crucial and specific to G_2.

5.1. **Parity decompositions of T and Y-systems.** For a triplet $(a, m, u) \in \mathcal{I}_\ell$, we reset the parity conditions $\mathbf{P}_+$ and $\mathbf{P}_-$ by

$$\mathbf{P}_+ : a + m + 3u \text{ is even,} \tag{5.1}$$

$$\mathbf{P}_- : a + m + 3u \text{ is odd.} \tag{5.2}$$

Then, we have $\mathcal{T}^\circ_\ell(G_2)_+ \simeq \mathcal{T}^\circ_\ell(G_2)_-$ by $T^{(a)}_m(u) \mapsto T^{(a)}_m(u + \frac{1}{3})$ and

$$\mathcal{T}^\circ_\ell(G_2) \simeq \mathcal{T}^\circ_\ell(G_2)_+ \otimes_\mathbb{Z} \mathcal{T}^\circ_\ell(G_2)_-. \tag{5.3}$$

For a triplet $(a, m, u) \in \mathcal{I}_\ell$, we reset the parity conditions $\mathbf{P}'_+$ and $\mathbf{P}'_-$ by

$$\mathbf{P}'_+ : a + m + 3u \text{ is odd,} \tag{5.4}$$

$$\mathbf{P}'_- : a + m + 3u \text{ is even.} \tag{5.5}$$



FIGURE 7. The quiver $Q_\ell(G_2)$ for even ℓ (upper) and for odd ℓ (lower), where we identify the right columns in the three quivers.

We have

$$(a,m,u):\mathbf{P}'_+ \iff (a,m,u\pm \tfrac{1}{t_a}):\mathbf{P}_+. \tag{5.6}$$

Also, we have $\mathcal{Y}^\circ_\ell(G_2)_+ \simeq \mathcal{Y}^\circ_\ell(G_2)_-$ by $Y^{(a)}_m(u)\mapsto Y^{(a)}_m(u+\frac{1}{3})$, $1+Y^{(a)}_m(u)\mapsto 1+Y^{(a)}_m(u+\frac{1}{3})$, and

$$\mathcal{Y}^\circ_\ell(G_2)\simeq \mathcal{Y}^\circ_\ell(G_2)_+\times \mathcal{Y}^\circ_\ell(G_2)_-. \tag{5.7}$$

5.2. **Quiver $Q_\ell(G_2)$.** With type G_2 and $\ell\geq 2$ we associate the quiver $Q_\ell(G_2)$ by Figure 7, where the right columns in the three quivers are identified. Also we assign the empty or filled circle $\circ/\bullet$ to each vertex; furthermore, we assign the sign $+/-$ to each vertex of property $\bullet$, and one of the numbers I,…, VI to each vertex of property $\circ$.

Let us choose the index set $\mathbf{I}$ of the vertices of $Q_\ell(G_2)$ so that $\mathbf{i}=(i,i')\in\mathbf{I}$ represents the vertex at the i'th row (from the bottom) of the left column in the

ith quiver (from the left) for $i = 1, 2, 3$, and the one of the right column in any quiver for $i = 4$. Thus, $i = 1, \dots, 4$, and $i' = 1, \dots, \ell-1$ if $i \neq 4$ and $i' = 1, \dots, 3\ell-1$ if $i = 4$.

For a permutation s of $\{1, 2, 3\}$, let $\boldsymbol{\nu}_s$ be the permutation of $\mathbf{I}$ such that $\boldsymbol{\nu}_s(i, i') = (s(i), i')$ for $i \neq 4$ and $(4, i')$ for $i = 4$. Let $\boldsymbol{\omega}$ be the involution acting on $\mathbf{I}$ by the up-down reflection. Let $\boldsymbol{\nu}_s(Q_\ell(G_2))$ and $\boldsymbol{\omega}(Q_\ell(G_2))$ denote the quivers induced from $Q_\ell(G_2)$ by $\boldsymbol{\nu}_s$ and $\boldsymbol{\omega}$, respectively.

Lemma 5.1. *Let $Q = Q_\ell(G_2)$.*

(i) We have a periodic sequence of mutations of quivers

$$\begin{aligned}
&Q \overset{\mu^\bullet_+\mu^\circ_{\mathrm{I}}}{\longleftrightarrow} \boldsymbol{\nu}_{(23)}(Q)^{\mathrm{op}} \overset{\mu^\bullet_-\mu^\circ_{\mathrm{II}}}{\longleftrightarrow} \boldsymbol{\nu}_{(312)}(Q) \overset{\mu^\bullet_+\mu^\circ_{\mathrm{III}}}{\longleftrightarrow} \boldsymbol{\nu}_{(13)}(Q)^{\mathrm{op}}\\
&\overset{\mu^\bullet_-\mu^\circ_{\mathrm{IV}}}{\longleftrightarrow} \boldsymbol{\nu}_{(231)}(Q) \overset{\mu^\bullet_+\mu^\circ_{\mathrm{V}}}{\longleftrightarrow} \boldsymbol{\nu}_{(12)}(Q)^{\mathrm{op}} \overset{\mu^\bullet_-\mu^\circ_{\mathrm{VI}}}{\longleftrightarrow} Q.
\end{aligned} \tag{5.8}$$

(ii) $\boldsymbol{\omega}(Q) = Q$ if $h^\vee + \ell$ is even, and $\boldsymbol{\omega}(Q) = \boldsymbol{\nu}_{(13)}(Q)^{\mathrm{op}}$ if $h^\vee + \ell$ is odd.

See Figures 8–10 for an example.

5.3. **Cluster algebra and alternative labels.**

Let $B_\ell(G_2)$ be the corresponding skew-symmetric matrix to the quiver $Q_\ell(G_2)$. In the rest of the section, we set the matrix $B = (B_{\mathbf{ij}})_{\mathbf{i},\mathbf{j}\in\mathbf{I}} = B_\ell(G_2)$ unless otherwise mentioned.

Let $\mathcal{A}(B, x, y)$ be the cluster algebra with coefficients in the universal semifield $\mathbb{Q}_{\mathrm{sf}}(y)$, and $\mathcal{G}(B, y)$ be the coefficient group associated with $\mathcal{A}(B, x, y)$.

In view of Lemma 5.1 we set $x(0) = x$, $y(0) = y$ and define clusters $x(u) = (x_{\mathbf{i}}(u))_{\mathbf{i}\in\mathbf{I}}$ $(u \in \frac{1}{3}\mathbb{Z})$ and coefficient tuples $y(u) = (y_{\mathbf{i}}(u))_{\mathbf{i}\in\mathbf{I}}$ $(u \in \frac{1}{3}\mathbb{Z})$ by the sequence of mutations

$$\begin{aligned}
&\cdots \overset{\mu^\bullet_-\mu^\circ_{\mathrm{VI}}}{\longleftrightarrow} (B, x(0), y(0)) \overset{\mu^\bullet_+\mu^\circ_{\mathrm{I}}}{\longleftrightarrow} (-\boldsymbol{\nu}_{(23)}(B), x(\tfrac{1}{3}), y(\tfrac{1}{3}))\\
&\overset{\mu^\bullet_-\mu^\circ_{\mathrm{II}}}{\longleftrightarrow} (\boldsymbol{\nu}_{(312)}(B), x(\tfrac{2}{3}), y(\tfrac{2}{3})) \overset{\mu^\bullet_+\mu^\circ_{\mathrm{III}}}{\longleftrightarrow} (-\boldsymbol{\nu}_{(13)}(B), x(1), y(1))\\
&\overset{\mu^\bullet_-\mu^\circ_{\mathrm{IV}}}{\longleftrightarrow} (\boldsymbol{\nu}_{(231)}(B), x(\tfrac{4}{3}), y(\tfrac{4}{3})) \overset{\mu^\bullet_+\mu^\circ_{\mathrm{V}}}{\longleftrightarrow} (-\boldsymbol{\nu}_{(12)}(B), x(\tfrac{5}{3}), y(\tfrac{5}{3})) \overset{\mu^\bullet_-\mu^\circ_{\mathrm{VI}}}{\longleftrightarrow} \cdots .
\end{aligned} \tag{5.9}$$

where $\boldsymbol{\nu}_s(B) = B'$ is defined by $B'_{\boldsymbol{\nu}_s(\mathbf{i})\boldsymbol{\nu}_s(\mathbf{j})} = B_{\mathbf{ij}}$.

For a pair $(\mathbf{i}, u) \in \mathbf{I} \times \frac{1}{3}\mathbb{Z}$, we set the parity condition $\mathbf{p}_+$ and $\mathbf{p}_-$ by

$$\mathbf{p}_+ : \begin{cases}
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{I}} \sqcup \mathbf{I}^\bullet_+ & u \equiv 0\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{II}} \sqcup \mathbf{I}^\bullet_- & u \equiv \frac{1}{3}\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{III}} \sqcup \mathbf{I}^\bullet_+ & u \equiv \frac{2}{3}\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{IV}} \sqcup \mathbf{I}^\bullet_- & u \equiv 1\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{V}} \sqcup \mathbf{I}^\bullet_+ & u \equiv \frac{4}{3}\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{VI}} \sqcup \mathbf{I}^\bullet_- & u \equiv \frac{5}{3},
\end{cases}
\qquad
\mathbf{p}_- : \begin{cases}
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{VI}} \sqcup \mathbf{I}^\bullet_- & u \equiv 0\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{I}} \sqcup \mathbf{I}^\bullet_+ & u \equiv \frac{1}{3}\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{II}} \sqcup \mathbf{I}^\bullet_- & u \equiv \frac{2}{3}\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{III}} \sqcup \mathbf{I}^\bullet_+ & u \equiv 1\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{IV}} \sqcup \mathbf{I}^\bullet_- & u \equiv \frac{4}{3}\\
\mathbf{i} \in \mathbf{I}^\circ_{\mathrm{V}} \sqcup \mathbf{I}^\bullet_+ & u \equiv \frac{5}{3},
\end{cases} \tag{5.10}$$

where $\equiv$ is modulo $2\mathbb{Z}$. We have

$$(\mathbf{i}, u) : \mathbf{p}_+ \iff (\mathbf{i}, u + \frac{1}{3}) : \mathbf{p}_-. \tag{5.11}$$

Each $(\mathbf{i}, u) : \mathbf{p}_+$ is a mutation point of (5.9) in the forward direction of u, and each $(\mathbf{i}, u) : \mathbf{p}_-$ is so in the backward direction of u.


$x^{(2)}_{11}(-\frac{1}{3})$
$x^{(2)}_{9}(-\frac{1}{3})$
$x^{(2)}_{7}(-\frac{1}{3})$
$x(0)$
$x^{(1)}_{2}(-1)$
$x^{(2)}_{5}(-\frac{1}{3})$
$x^{(2)}_{3}(-\frac{1}{3})$
$x^{(2)}_{1}(-\frac{1}{3})$
$\mu^{\bullet}_{+}\mu^{\circ}_{\mathrm{I}} \updownarrow$
$x^{(2)}_{10}(0)$
$x^{(1)}_{3}(-\frac{2}{3})$
$x^{(2)}_{8}(0)$
$x(\frac{1}{3})$
$x^{(2)}_{6}(0)$
$x^{(2)}_{4}(0)$
$x^{(1)}_{1}(-\frac{2}{3})$
$x^{(2)}_{2}(0)$
$\mu^{\bullet}_{-}\mu^{\circ}_{\mathrm{II}} \updownarrow$

FIGURE 8. (Continues to Figures 9 and 10) Label of cluster variables $x_{\mathbf{i}}(u)$ by $\mathcal{I}_{\ell+}$ for G_2, $\ell = 4$. The right columns in the middle and right quivers (marked by $\diamond$) are identified with the right column in the left quiver.


$x^{(2)}_{11}(\frac{1}{3})$
$x^{(2)}_{9}(\frac{1}{3})$
$x^{(2)}_{7}(\frac{1}{3})$
$x(\frac{2}{3})$
$x^{(1)}_{2}(-\frac{1}{3})$
$x^{(2)}_{5}(\frac{1}{3})$
$x^{(2)}_{3}(\frac{1}{3})$
$x^{(2)}_{1}(\frac{1}{3})$
$\mu^{\bullet}_{+}\mu^{\circ}_{\mathrm{III}}\updownarrow$
$x^{(2)}_{10}(\frac{2}{3})$
$x^{(1)}_{3}(0)$
$x^{(2)}_{8}(\frac{2}{3})$
$x(1)$
$x^{(2)}_{6}(\frac{2}{3})$
$x^{(2)}_{4}(\frac{2}{3})$
$x^{(1)}_{1}(0)$
$x^{(2)}_{2}(\frac{2}{3})$
$\mu^{\bullet}_{-}\mu^{\circ}_{\mathrm{IV}}\updownarrow$

FIGURE 9. (Continues from Figure 8)


$x^{(2)}_{11}(1)$
$x^{(2)}_{9}(1)$
$x^{(2)}_{7}(1)$
$x(\frac{4}{3})$
$x^{(1)}_{2}(\frac{1}{3})$
$x^{(2)}_{5}(1)$
$x^{(2)}_{3}(1)$
$x^{(2)}_{1}(1)$
$\mu^{\bullet}_{+}\mu^{\circ}_{\mathrm{V}}\updownarrow$
$x^{(2)}_{10}(\frac{4}{3})$
$x^{(1)}_{3}(\frac{2}{3})$
$x^{(2)}_{8}(\frac{4}{3})$
$x(\frac{5}{3})$
$x^{(2)}_{6}(\frac{4}{3})$
$x^{(2)}_{4}(\frac{4}{3})$
$x^{(1)}_{1}(\frac{2}{3})$
$x^{(2)}_{2}(\frac{4}{3})$
$\mu^{\bullet}_{-}\mu^{\circ}_{\mathrm{VI}}\updownarrow$

FIGURE 10. (Continues from Figure 9)

Lemma 5.2. *Below $\equiv$ means the equivalence modulo $2\mathbb{Z}$.*

(i) The map

$$\begin{array}{rccl} g: & \mathcal{I}_{\ell+} & \to & \{(\mathbf{i},u):\mathbf{p}_+\} \\ & (a,m,u-\frac{1}{t_a}) & \mapsto & \begin{cases} ((1,m),u) & a=1;\, m+u\equiv 0 \\ ((2,m),u) & a=1;\, m+u\equiv \frac{4}{3} \\ ((3,m),u) & a=1;\, m+u\equiv \frac{2}{3} \\ ((4,m),u) & a=2 \end{cases} \end{array} \tag{5.12}$$

is a bijection.

(ii) The map

$$\begin{array}{rccl} g': & \mathcal{I}'_{\ell+} & \to & \{(\mathbf{i},u):\mathbf{p}_+\} \\ & (a,m,u) & \mapsto & \begin{cases} ((1,m),u) & a=1;\, m+u\equiv 0 \\ ((2,m),u) & a=1;\, m+u\equiv \frac{4}{3} \\ ((3,m),u) & a=1;\, m+u\equiv \frac{2}{3} \\ ((4,m),u) & a=2 \end{cases} \end{array} \tag{5.13}$$

is a bijection.

We introduce alternative labels $x_{\mathbf{i}}(u)=x^{(a)}_m(u-1/t_a)$ $((a,m,u-1/t_a)\in\mathcal{I}_{\ell+})$ for $(\mathbf{i},u)=g((a,m,u-1/t_a))$ and $y_{\mathbf{i}}(u)=y^{(a)}_m(u)$ $((a,m,u)\in\mathcal{I}'_{\ell+})$ for $(\mathbf{i},u)=g'((a,m,u))$, respectively. See Figures 8–10 for an example.

5.4. **T-system and cluster algebra.**

Lemma 5.3. *The family $\{x^{(a)}_m(u)\mid(a,m,u)\in\mathcal{I}_{\ell+}\}$ satisfies the system of relations* (3.16) *with $G(b,k,v;a,m,u)$ for $\mathbb{T}_\ell(G_2)$. In particular, the family $\{[x^{(a)}_m(u)]_{\mathbf{1}}\mid(a,m,u)\in\mathcal{I}_{\ell+}\}$ satisfies the T-system $\mathbb{T}_\ell(G_2)$ in $\mathcal{A}(B,x)$ by replacing $T^{(a)}_m(u)$ with $[x^{(a)}_m(u)]_{\mathbf{1}}$.*

Definition 5.4. The *T-subalgebra $\mathcal{A}_T(B,x)$ of $\mathcal{A}(B,x)$ associated with the sequence* (5.9) is the subring of $\mathcal{A}(B,x)$ generated by $[x_{\mathbf{i}}(u)]_{\mathbf{1}}$ $((\mathbf{i},u)\in\mathbf{I}\times\frac{1}{3}\mathbb{Z})$.

Theorem 5.5. *The ring $\mathcal{T}^\circ_\ell(G_2)_+$ is isomorphic to $\mathcal{A}_T(B,x)$ by the correspondence $T^{(a)}_m(u)\mapsto[x^{(a)}_m(u)]_{\mathbf{1}}$.*

5.5. **Y-system and cluster algebra.**

Lemma 5.6. *The family $\{y^{(a)}_m(u)\mid(a,m,u)\in\mathcal{I}'_{\ell+}\}$ satisfies the Y-system $\mathbb{Y}_\ell(G_2)$ by replacing $Y^{(a)}_m(u)$ with $y^{(a)}_m(u)$.*

Definition 5.7. The *Y-subgroup $\mathcal{G}_Y(B,y)$ of $\mathcal{G}(B,y)$ associated with the sequence* (5.9) is the subgroup of $\mathcal{G}(B,y)$ generated by $y_{\mathbf{i}}(u)$ $((\mathbf{i},u)\in\mathbf{I}\times\frac{1}{3}\mathbb{Z})$ and $1+y_{\mathbf{i}}(u)$ $((\mathbf{i},u):\mathbf{p}_+$ or $\mathbf{p}_-)$.

Theorem 5.8. *The group $\mathcal{Y}^\circ_\ell(G_2)_+$ is isomorphic to $\mathcal{G}_Y(B,y)$ by the correspondence $Y^{(a)}_m(u)\mapsto y^{(a)}_m(u)$ and $1+Y^{(a)}_m(u)\mapsto 1+y^{(a)}_m(u)$.*

5.6. **Tropical Y-system at level 2.** By direct computations, the following properties are verified.

Proposition 5.9. *For* $[\mathcal{G}_Y(B,y)]_{\mathbf{T}}$ *with* $B = B_2(G_2)$*, the following facts hold.*

(i) Let u *be in the region* $0 \leq u < 2$*. For any* $(\mathbf{i}, u) : \mathbf{p}_+$*, the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is positive.*

(ii) Let u *be in the region* $-h^\vee \leq u < 0$*.*

(a) Let $\mathbf{i} = (1,1), (2,1), (3,1)$*, or* $(4,3)$*. For any* $(\mathbf{i}, u) : \mathbf{p}_+$*, the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is negative.*

(b) Let $\mathbf{i} = (4,1), (4,2), (4,4)$*, or* $(4,5)$ *For any* $(\mathbf{i}, u) : \mathbf{p}_+$*, the monomial* $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ *is negative for* $u = -\frac{1}{3}, -\frac{2}{3}, -2, -\frac{7}{3}, -\frac{11}{3}, -4$ *and positive for* $u = -1, -\frac{4}{3}, -\frac{5}{3}, -\frac{8}{3}, -3, -\frac{10}{3}$*.*

(iii) $y_{ii'}(2) = y_{ii'}^{-1}$ *if* $i \neq 4$ *and* $y_{4,6-i'}^{-1}$ *if* $i = 4$*.*

(iv) $y_{ii'}(-h^\vee) = y_{ii'}^{-1}$*.*

Also we have a description of the core part of $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ in the region $-h^\vee \leq u < 0$ in terms of the root system of D_4. We use the following index of the Dynkin diagram D_4.



Let $\Pi = \{\alpha_1, \dots, \alpha_4\}$, $-\Pi$, Φ_+ be the set of the simple roots, the negative simple roots, the positive roots, respectively, of type D_4. Let σ_i be the piecewise-linear analogue of the simple reflection s_i, acting on the set of the almost positive roots $\Phi_{\geq -1} = \Phi_+ \sqcup (-\Pi)$. We define σ as the composition

$$\sigma = \sigma_3\sigma_4\sigma_1\sigma_4\sigma_2\sigma_4. \tag{5.14}$$

Lemma 5.10. *We have the orbits by* σ

$$\begin{aligned}
&-\alpha_1 \to \alpha_1 + \alpha_3 + \alpha_4 \to \alpha_1 + \alpha_2 + \alpha_4 \to -\alpha_1,\\
&-\alpha_2 \to \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 \to \alpha_2 + \alpha_4 \to -\alpha_2,\\
&-\alpha_3 \to \alpha_3 \to \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 \to -\alpha_3,\\
&-\alpha_4 \to \alpha_2 + \alpha_3 + \alpha_4 \to \alpha_4 \to \alpha_1\\
&\qquad \to \alpha_2 \to \alpha_3 + \alpha_4 \to \alpha_1 + \alpha_4 \to -\alpha_4.
\end{aligned} \tag{5.15}$$

In particular, these elements in Φ_+ *exhaust the set* Φ_+*, thereby providing the orbit decomposition of* Φ_+ *by* σ*.*

For $-h^\vee \leq u < 0$, define

$$\alpha_i(u) = \begin{cases}
\sigma^{-(u-1)/2}(-\alpha_1) & i = 1;\ u = -1, -3,\\
\sigma^{-(3u-1)/6}(-\alpha_2) & i = 2;\ u = -\frac{5}{3}, -\frac{11}{3},\\
\sigma^{-(3u-5)/6}(-\alpha_3) & i = 3;\ u = -\frac{1}{3}, -\frac{7}{3},\\
\sigma^{-(3u+2)/6}(\alpha_3 + \alpha_4) & i = 4;\ u = -\frac{2}{3}, -\frac{8}{3},\\
\sigma^{-(3u+4)/6}(\alpha_1) & i = 4;\ u = -\frac{4}{3}, -\frac{10}{3},\\
\sigma^{-u/2}(-\alpha_4) & i = 4;\ u = -2, -4.
\end{cases} \tag{5.16}$$

By Lemma 5.10 they are (all the) positive roots of D_4.

For a monomial m in $y = (y_{\mathbf{i}})_{\mathbf{i}\in\mathbf{I}}$, let $\pi_D(m)$ denote the specialization with $y_{41} = y_{42} = y_{44} = y_{45} = 1$. For simplicity, we set $y_{i1} = y_i$ $(i \neq 4)$, $y_{43} = y_4$, and also, $y_{i1}(u) = y_i(u)$ $(i \neq 4)$, $y_{43}(u) = y_4(u)$. We define the vectors $\mathbf{t}_i(u) = (t_i(u)_k)_{k=1}^4$ by

$$\pi_D([y_i(u)]_{\mathbf{T}}) = \prod_{k=1}^{4} y_k^{t_i(u)_k}. \tag{5.17}$$

We also identify each vector $\mathbf{t}_i(u)$ with $\alpha = \sum_{k=1}^4 t_i(u)_k\alpha_k \in \mathbb{Z}\Pi$.

Proposition 5.11. *Let $-h^\vee \leq u < 0$. Then, we have*

$$\mathbf{t}_i(u) = -\alpha_i(u), \tag{5.18}$$

for (i, u) in (5.16).

5.7. **Tropical Y-systems of higher levels.**

Proposition 5.12. *Let $\ell > 2$ be an integer. For $[\mathcal{G}_Y(B, y)]_{\mathbf{T}}$ with $B = B_\ell(G_2)$, the following facts hold.*

(i) Let u be in the region $0 \leq u < \ell$. For any $(\mathbf{i}, u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is positive.

(ii) Let u be in the region $-h^\vee \leq u < 0$.

- *(a) Let $\mathbf{i} \in \mathbf{I}^\circ$ or $(4, i')$ $(i' \in 3\mathbb{N})$. For any $(\mathbf{i}, u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is negative.*
- *(b) Let $\mathbf{i} = (4, i')$ $(i' \notin 3\mathbb{N})$. For any $(\mathbf{i}, u) : \mathbf{p}_+$, the monomial $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ is negative for $u = -\frac{1}{3}, -\frac{2}{3}, -2, -\frac{7}{3}, -\frac{11}{3}, -4$ and positive for $u = -1, -\frac{4}{3}, -\frac{5}{3}, -\frac{8}{3}, -3, -\frac{10}{3}$.*

(iii) $y_{ii'}(\ell) = y_{i,\ell-i'}^{-1}$ if $i \neq 4$ and $y_{4,3\ell-i'}^{-1}$ if $i = 4$.

(iv) $y_{ii'}(-h^\vee) = y_{ii'}^{-1}$.

We obtain corollaries of Propositions 5.9 and 5.12.

Theorem 5.13. *For $[\mathcal{G}_Y(B, y)]_{\mathbf{T}}$, the following relations hold.*

(i) Half periodicity: $[y_{\mathbf{i}}(u + h^\vee + \ell)]_{\mathbf{T}} = [y_{\boldsymbol{\omega}(\mathbf{i})}(u)]_{\mathbf{T}}$.

(ii) Full periodicity: $[y_{\mathbf{i}}(u + 2(h^\vee + \ell))]_{\mathbf{T}} = [y_{\mathbf{i}}(u)]_{\mathbf{T}}$.

Theorem 5.14. *For $[\mathcal{G}_Y(B, y)]_{\mathbf{T}}$, let N_+ and N_- denote the total numbers of the positive and negative monomials, respectively, among $[y_{\mathbf{i}}(u)]_{\mathbf{T}}$ for $(\mathbf{i}, u) : \mathbf{p}_+$ in the region $0 \leq u < 2(h^\vee + \ell)$. Then, we have*

$$N_+ = 6\ell(2\ell + 1), \quad N_- = 12(3\ell - 2). \tag{5.19}$$

5.8. **Periodicities and dilogarithm identities.** Applying [IIKKN, Theorem 5.1] to Theorem 5.13, we obtain the periodicities:

Theorem 5.15. *For $\mathcal{A}(B, x, y)$, the following relations hold.*

(i) Half periodicity: $x_{\mathbf{i}}(u + h^\vee + \ell) = x_{\boldsymbol{\omega}(\mathbf{i})}(u)$.

(ii) Full periodicity: $x_{\mathbf{i}}(u + 2(h^\vee + \ell)) = x_{\mathbf{i}}(u)$.

Theorem 5.16. *For $\mathcal{G}(B, y)$, the following relations hold.*

(i) Half periodicity: $y_{\mathbf{i}}(u + h^\vee + \ell) = y_{\boldsymbol{\omega}(\mathbf{i})}(u)$.

(ii) Full periodicity: $y_{\mathbf{i}}(u + 2(h^\vee + \ell)) = y_{\mathbf{i}}(u)$.

Then, Theorems 2.5 and 2.6 for G_2 follow from Theorems 5.5, 5.8, 5.15, and 5.16. Furthermore, Theorem 2.10 for G_2 is obtained from the above periodicities and Theorem 5.14 as in the B_r case [IIKKN, Section 6].

6. Mutation equivalence of quivers

Recall that two quivers Q and Q' are said to be *mutation equivalent*, and denoted by $Q \sim Q'$ here, if there is a quiver isomorphism from Q to some quiver obtained from Q' by successive mutations.

Below we present several mutation equivalent pairs of the quivers $Q_\ell(X_r)$, though the list is not complete at all. For simply laced X_r, $Q_\ell(X_r)$ is the quiver defined as the square product $\vec{X}_r \square \vec{A}_{\ell-1}$ in [Ke, Section 8].

Proposition 6.1. *We have the following mutation equivalences of quivers.*

$$\begin{aligned} Q_2(B_r) &\sim Q_2(D_{2r+1}), \\ Q_2(C_3) &\sim Q_3(D_4), \\ Q_2(F_4) &\sim Q_3(D_5), \\ Q_3(C_2) &\sim Q_4(A_3), \\ Q_\ell(G_2) &\sim Q_\ell(C_3). \end{aligned} \tag{6.1}$$

References

[FZ1] S. Fomin, A. Zelevinsky, Y-systems and generalized associahedra, *Ann. of Math.* **158** (2003), 977–1018.

[FZ2] S. Fomin, A. Zelevinsky, Cluster algebras IV. Coefficients, *Compositio Mathematica* **143** (2007), 112–164.

[IIKKN] R. Inoue, O. Iyama, B. Keller, A. Kuniba, T. Nakanishi, Periodicities of T and Y-systems, dilogarithm identities, and cluster algebras I: Type B_r, arXiv:1001.1880.

[IIKNS] R. Inoue, O. Iyama, A. Kuniba, T. Nakanishi, J. Suzuki, Periodicities of T-systems and Y-systems, *Nagoya Math. J.* **197** (2010), 59–174.

[Ke] B. Keller, Cluster algebras, quiver representations and triangulated categories, in *Triangulated categories*, T. Holm, P. Jørgensen, and R. Rouquier, eds., London Mathematical Society, Lecture Note Series vol. 375, Cambridge University Press, 2010, pp. 76–160.

[Ki] A. N. Kirillov, Identities for the Rogers dilogarithm function connected with simple Lie algebras, *J. Sov. Math.* **47** (1989), 2450–2458.

[Ku] A. Kuniba, Thermodynamics of the $U_q(X_r^{(1)})$ Bethe ansatz system with q a root of unity, *Nucl. Phys.* **B389** (1993), 209–244.

[KN] A. Kuniba, T. Nakanishi, Spectra in conformal field theories from the Rogers dilogarithm, *Mod. Phys. Lett.* **A7** (1992), 3487–3494.

[KNS] A. Kuniba, T. Nakanishi, J. Suzuki, Functional relations in solvable lattice models: I. Functional relations and representation theory, *Int. J. Mod. Phys. A* **9** (1994), 5215–5266.

[P1] P. Plamondon, Cluster characters for cluster categories with infinite-dimensional morphism spaces, arXiv:1002.4956.

[P2] P. Plamondon, Cluster algebras via cluster categories with infinite-dimensional morphism spaces, arXiv:1004.0830.

R. Inoue: Faculty of Pharmaceutical Sciences, Suzuka University of Medical Science, Suzuka, 513-8670, Japan

O. Iyama: Graduate School of Mathematics, Nagoya University, Nagoya, 464-8604, Japan

B. Keller: Université Paris Diderot – Paris 7, UFR de Mathématiques, Institut de Mathématiques de Jussieu, UMR 7586 du CNRS, Case 7012, 2, place Jussieu, 75251 Paris Cedex 05, France

A. Kuniba: Institute of Physics, University of Tokyo, Tokyo, 153-8902, Japan

T. Nakanishi: Graduate School of Mathematics, Nagoya University, Nagoya, 464-8604, Japan